

【High Fat Diet 32(HFD32)飼料参考文献リスト】

2005

Oike Y, Akao M, Yasunaga K, Yamauchi T, Morisada T, Ito Y, Urano T, Kimura Y, Kubota Y, Maekawa H, Miyamoto T, Miyata K, Matsumoto S, Sakai J, Nakagata N, Takeya M, Koseki H, Ogawa Y, Kadokawa T, Suda T. Angiopoietin-related growth factor antagonizes obesity and insulin resistance. *Nat Med.* 2005; 11(4): 400-8. [PMID: 15778720]

2007

Nasu R, Seki K, Nara M, Murakami M, Kohama T. Effect of a high-fat diet on diabetic mother rats and their offspring through three generations. *Endocr J.* 2007; 54(4): 563-9. [PMID: 17603226]

Yamato M, Shiba T, Yoshida M, Ide T, Seri N, Kudou W, Kinugawa S, Tsutsui H. Fatty acids increase the circulating levels of oxidative stress factors in mice with diet-induced obesity via redox changes of albumin. *FEBS J.* 2007; 274(15): 3855-63. [PMID: 17617229]

Nishikawa S, Yasoshima A, Doi K, Nakayama H, Uetsuka K. Involvement of sex, strain and age factors in high fat diet-induced obesity in C57BL/6J and BALB/cA mice. *Exp Anim.* 2007; 56(4): 263-72. [PMID: 17660680]

Kamei Y, Suganami T, Kohda T, Ishino F, Yasuda K, Miura S, Ezaki O, Ogawa Y. Peg1/Mest in obese adipose tissue is expressed from the paternal allele in an isoform-specific manner. *FEBS Lett.* 2007; 581(1): 91-6. [PMID: 17182038]

Aoyagi T, Birumachi J, Hiroyama M, Fujiwara Y, Sanbe A, Yamauchi J, Tanoue A. Alteration of glucose homeostasis in V1a vasopressin receptor-deficient mice. *Endocrinology.* 2007; 148(5): 2075-84. [PMID: 17303660]

Nara N, Nakayama Y, Okamoto S, Tamura H, Kiyono M, Muraoka M, Tanaka K, Taya C, Shitara H, Ishii R, Yonekawa H, Minokoshi Y, Hara T. Disruption of CXCR Motif Chemokine Ligand-14 in Mice Ameliorates Obesity-induced Insulin Resistance. *J Biol Chem.* 2007; 282(42): 30794-803. [PMID: 17724031]

2008

Sakurai T, Nishioka H, Fujii H, Nakano N, Kizaki T, Radak Z, Izawa T, Haga S, Ohno H. Antioxidative effects of a new lychee fruit-derived polyphenol mixture, oligonol, converted into a low-molecular form in adipocytes. *Biosci Biotechnol Biochem.* 2008; 72(2): 463-76. [PMID: 18256485]

Nishikawa S, Doi K, Nakayama H, Uetsuka K. The effect of fasting on hepatic lipid accumulation and transcriptional regulation of lipid metabolism differs between C57BL/6J and BALB/cA mice fed a high-fat diet. *Toxicol Pathol.* 2008; 36(6): 850-7. [PMID: 18812581]

Kobayashi M, Ohnishi H, Okazawa H, Murata Y, Hayashi Y, Kobayashi H, Kitamura T, Matozaki T. Expression of Src homology 2 domain-containing protein tyrosine phosphatase substrate-1 in pancreatic beta-cells and its role in promotion of insulin secretion and protection against diabetes. *Endocrinology.* 2008; 149(11): 5662-9. [PMID: 18635660]

2009

Ohori H, Yamakoshi H, Tomizawa M, Shibuya M, Kakudo Y, Takahashi A, Takahashi S, Kato S, Suzuki T, Ishioka C, Iwabuchi Y, Shibata H. Synthesis and biological analysis of new curcumin analogues bearing an enhanced potential for the medicinal treatment of cancer. *Mol Cancer Ther.* 2006; 5(10): 2563-71. [PMID: 17041101]

2009

Horii T, Morita S, Kimura M, Hatada I. Epigenetic regulation of adipocyte differentiation by a Rho guanine nucleotide exchange factor, WGEF. *PLoS One.* 2009; 4(6): e5809. [PMID: 19503838]

Fujimoto T, Miyasaka K, Koyanagi M, Tsunoda T, Baba I, Doi K, Ohta M, Kato N, Sasazuki T, Shirasawa S. Altered energy homeostasis and resistance to diet-induced obesity in KRAP-deficient mice. *PLoS One.* 2009; 4(1): e4240. [PMID: 19156225]

Kubo M, Ijichi N, Ikeda K, Horie-Inoue K, Takeda S, Inoue S. Modulation of adipogenesis-related gene expression by estrogen-related receptor gamma during adipocytic differentiation. *Biochim Biophys Acta.* 2009; 1789(2): 71-7. [PMID: 18809516]

Tozuka Y, Wada E, Wada K. Diet-induced obesity in female mice leads to peroxidized lipid accumulations and impairment of hippocampal neurogenesis during the early life of their offspring. *FASEB J.* 2009; 23(6): 1920-34. [PMID: 19158155]2009

Nakamura K, Aoyagi T, Hiroyama M, Kusakawa S, Mizutani R, Sanbe A, Yamauchi J, Kamohara M, Momose K, Tanoue A. Both V(1A) and V(1B) vasopressin receptors deficiency result in impaired glucose tolerance. *Eur J Pharmacol.*

2009; 613(1-3): 182-8. [PMID: 19375419]

Matsuhashima S, Kinugawa S, Yokota T, Inoue N, Ohta Y, Hamaguchi S, Tsutsui H. Increased myocardial NAD(P)H oxidase-derived superoxide causes the exacerbation of postinfarct heart failure in type 2 diabetes. *Am J Physiol Heart Circ Physiol.* 2009; 297(1): H409-16. [PMID: 19465539]

Nozaki Y, Fujita K, Yoneda M, Wada K, Shinohara Y, Takahashi H, Kirikoshi H, Inamori M, Kubota K, Saito S, Mizoue T, Masaki N, Nagashima Y, Terauchi Y, Nakajima A. Long-term combination therapy of ezetimibe and acarbose for non-alcoholic fatty liver disease. *J Hepatol.* 2009; 51(3): 548-56. [PMID: 19596472]

Yokota T, Kinugawa S, Hirabayashi K, Matsuhashima S, Inoue N, Ohta Y, Hamaguchi S, Sobrin MA, Ono T, Suga T, Kuroda S, Tanaka S, Terasaki F, Okita K, Tsutsui H. Oxidative stress in skeletal muscle impairs mitochondrial respiration and limits exercise capacity in type 2 diabetic mice. *Am J Physiol Heart Circ Physiol.* 2009; 297(3): H1069-77. [PMID: 19617406]

Matsubara Y, Kano K, Kondo D, Mugishima H, Matsumoto T: Differences in Adipocytokines and Fatty Acid Composition between Two Adipocyte Fractions of Small and Large Cells in High-Fat Diet-Induced Obese Mice. *Ann Nutr Metab* 2009; 54: 258-267 [PMID: 19641303]

Endo H, Hosono K, Fujisawa T, Takahashi H, Sugiyama M, Yoneda K, Nozaki Y, Fujita K, Yoneda M, Inamori M, Wada K, Nakagama H, Nakajima A. Involvement of JNK pathway in the promotion of the early stage of colorectal carcinogenesis under high-fat dietary conditions. *Gut.* 2009; 58(12): 1637-43. [PMID: 19570763]

Shibata H, Yamakoshi H, Sato A, Ohori H, Kakudo Y, Kudo C, Takahashi Y, Watanabe M, Takano H, Ishioka C, Noda T, Iwabuchi Y. Newly synthesized curcumin analog has improved potential to prevent colorectal carcinogenesis in vivo. *Cancer Sci.* 2009; 100(5): 956-60. [PMID: 19445025]

Setsuie R, Suzuki M, Kabuta T, Fujita H, Miura S, Ichihara N, Yamada D, Wang YL, Ezaki O, Suzuki Y, Wada K. Ubiquitin C-terminal hydrolase-L3-knockout mice are resistant to diet-induced obesity and show increased activation of AMP-activated protein kinase in skeletal muscle. *FASEB J.* 2009; 23(12): 4148-57. [PMID: 19671667]

Tabata M, Kadomatsu T, Fukuhara S, Miyata K, Ito Y, Endo M, Urano T, Zhu HJ, Tsukano H, Tazume H, Kaikita K, Miyashita K, Iwawaki T, Shimabukuro M, Sakaguchi K, Ito T, Nakagata N, Yamada T, Katagiri H, Kasuga M, Ando Y, Ogawa H, Mochizuki N, Itoh H, Suda T, Oike Y.

Angiopoietin-like protein 2 promotes chronic adipose tissue inflammation and obesity-related systemic insulin resistance. *Cell Metab.* 2009; 10(3): 178-88. [PMID: 19723494]

Shimoda H, Tanaka J, Kikuchi M, Fukuda T, Ito H, Hatano T, Yoshida T. Effect of polyphenol-rich extract from walnut on diet-induced hypertriglyceridemia in mice via enhancement of fatty acid oxidation in the liver. *J Agric Food Chem.* 2009; 57(5): 1786-92. [PMID: 19256553]

2010

Hasegawa G, Yamasaki M, Kadono M, Tanaka M, Asano M, Senmaru T, Kondo Y, Fukui M, Obayashi H, Maruyama N, Nakamura N, Ishigami A. Senescence marker protein-30/gluconolactonase deletion worsens glucose tolerance through impairment of acute insulin secretion. *Endocrinology.* 2010; 151(2): 529-36. [PMID: 19934374]

Tanegashima K, Suzuki K, Nakayama Y, Hara T. Antibody-assisted enhancement of biological activities of CXCL14 in human monocytic leukemia-derived THP-1 cells and high fat diet-induced obese mice. *Exp Cell Res.* 2010; 316(7): 1263-70. [PMID: 20083103]

Tozuka Y, Kumon M, Wada E, Onodera M, Mochizuki H, Wada K. Maternal obesity impairs hippocampal BDNF production and spatial learning performance in young mouse offspring. *Neurochem Int.* 2010; 57(3): 235-47. [PMID: 20538025]

Miyagi T, Takehara T, Uemura A, Nishio K, Shimizu S, Kodama T, Hikita H, Li W, Sasakawa A, Tatsumi T, Ohkawa K, Kanto T, Hiramatsu N, Hayashi N. Absence of invariant natural killer T cells deteriorates liver inflammation and fibrosis in mice fed high-fat diet. *J Gastroenterol.* 2010; 45(12): 1247-54. [PMID: 20596733]

2011

Oiso H, Furukawa N, Suefuji M, Shimoda S, Ito A, Furumai R, Nakagawa J, Yoshida M, Nishino N, Araki E. The role of class I histone deacetylase (HDAC) on gluconeogenesis in liver. *Biochem Biophys Res Commun.* 2011; 404(1): 166-72. [PMID: 21108932]

Nakatani M, Kokubo M, Ohsawa Y, Sunada Y, Tsuchida K. Folistatin-derived peptide expression in muscle decreases adipose tissue mass and prevents hepatic steatosis. *Am J Physiol Endocrinol Metab.* 2011; 300(3): E543-53. [PMID: 21205933]

Kuwashiro S, Terai S, Oishi T, Fujisawa K, Matsumoto T, Nishina H, Sakaieda I. Telmisartan improves nonalcoholic steatohepatitis in medaka (*Oryzias latipes*) by reducing

- macrophage infiltration and fat accumulation. *Cell Tissue Res.* 2011; 344(1): 125-34. [PMID: 21327395]
- Tominaga K, Yamauchi A, Egawa T, Tanaka R, Kawahara S, Shuto H, Kataoka Y. Vascular dysfunction and impaired insulin signaling in high-fat diet fed ovariectomized mice. *Microvasc Res.* 2011; 82(2): 171-6. [PMID: 21723877]
- Maejima Y, Iwasaki Y, Yamahara Y, Kodaira M, Sedbazar U, Yada T. Peripheral oxytocin treatment ameliorates obesity by reducing food intake and visceral fat mass. *Aging.* 2011; 3(12): 1169-77. [PMID: 22184277]
- Morita M, Oike Y, Nagashima T, Kadomatsu T, Tabata M, Suzuki T, Nakamura T, Yoshida N, Okada M, Yamamoto T. Obesity resistance and increased hepatic expression of catabolism-related mRNAs in Cnot3^{+/−} mice. *EMBO J.* 2011; 30(22): 4678-91. [PMID: 2189736]
- 大友理宣,高嶋亜希子,菊池継夫,高橋純一郎,戸枝一喜,畠恵司. 米糠乳酸発酵抽出物の高脂肪食負荷ラットにおける脂質異常改善作用. *生薬学雑誌.* 2011; 65: 33-8.
- Shirakawa J, Amo K, Ohminami H, Orime K, Togashi Y, Ito Y, Tajima K, Koganei M, Sasaki H, Takeda E, Terauchi Y. Protective effects of dipeptidyl peptidase-4 (DPP-4) inhibitor against increased β cell apoptosis induced by dietary sucrose and linoleic acid in mice with diabetes. *J Biol Chem.* 2011; 86(29): 25467-76. [PMID: 21613229]
- Tajima A, Hirata T, Taniguchi K, Kondo Y, Kato S, Saito-Hori M, Ishimoto T, Yamamoto K. Combination of TS-021 with metformin improves hyperglycemia and synergistically increases pancreatic β-cell mass in a mouse model of type 2 diabetes. *Life Sci.* 2011; 89(17-18): 662-70. [PMID: 21872612]
- Morita M, Oike Y, Nagashima T, Kadomatsu T, Tabata M, Suzuki T, Nakamura T, Yoshida N, Okada M, Yamamoto T. Obesity resistance and increased hepatic expression of catabolism-related mRNAs in Cnot3^{+/−} mice. *EMBO J.* 2011; 30(22): 4678-91. [PMID: 21897366]

2012

Yamato M, Shiba T, Ide T, Seri N, Kudo W, Ando M, Yamada K, Kinugawa S, Tsutsui H. High-fat diet-induced obesity and insulin resistance were ameliorated via enhanced fecal bile acid excretion in tumor necrosis factor-alpha receptor knockout mice. *Mol Cell Biochem.* 2012; 359(1-2): 161-7. [PMID: 21850461]

Febriza Dwiranti, Masanori Hiraoka, Takahiro Taguchi, Yuko Konishi, Mari Tominaga, Akira Tominaga. Effects of Gametophytes of *Ecklonia Kurome* on the Levels of Glucose and Triacylglycerol in db/db, Prediabetic C57BL/6J and IFN-γ KO Mice. *Int J B Sci.* 2012; 8(1): 64-75.

Satoh M, Andoh Y, Clingan CS, Ogura H, Fujii S, Eshima K, Nakayama T, Taniguchi M, Hirata N, Ishimori N, Tsutsui H, Onoé K, Iwabuchi K. Type II NKT cells stimulate diet-induced obesity by mediating adipose tissue inflammation, steatohepatitis and insulin resistance. *PLoS One.* 2012; 7(2): e30568. [PMID: 22383967]

Inazuka F, Sugiyama N, Tomita M, Abe T, Shioi G, Esumi H. Muscle-specific knock-out of NUAK family SNF1-like kinase 1 (NUAK1) prevents high fat diet-induced glucose intolerance. *J Biol Chem.* 2012; 287(20): 16379-89. [PMID: 22418434]

Sakurai Y, Inoue H, Shintani N, Arimori A, Hamagami KI, Hayata-Takano A, Baba A, Hashimoto H. Compensatory Recovery of Blood Glucose Levels in KKA(y) Mice Fed a High-Fat Diet: Insulin-Sparing Effects of PACAP Overexpression in β Cells. *J Mol Neurosci.* 2012; 48(3): 647-53. [PMID: 22477644]

Kawai D, Takaki A, Nakatsuka A, Wada J, Tamaki N, Yasunaka T, Koike K, Tsuzaki R, Matsumoto K, Miyake Y, Shiraha H, Morita M, Makino H, Yamamoto K. Hydrogen-rich water prevents progression of nonalcoholic steatohepatitis and accompanying hepatocarcinogenesis in mice. *Hepatology.* 2012; 56(3): 912-21. [PMID: 22505328]

Yamagata H, Ikejima K, Takeda K, Aoyama T, Kon K, Okumura K, Watanabe S. Altered expression and function of hepatic natural killer T cells in obese and diabetic KK-A(y) mice. *Hepatol Res.* 2012 Jul 19. [Epub ahead of print] [PMID: 22834991]

Nishikawa S, Sugimoto J, Okada M, Sakairi T, Takagi S. Gene expression in livers of BALB/C and C57BL/6J mice fed a high-fat diet. *Toxicol Pathol.* 2012; 40(1): 71-82. [PMID: 22105644]

Aizawa N, Homma Y, Igawa Y. Influence of High Fat Diet Feeding for 20 Weeks on Lower Urinary Tract Function in Mice. LUTS: Lower Urinary Tract Symptoms Early View, Article first published online: 21 SEP 2012.

Burdeos GC, Nakagawa K, Kimura F, Miyazawa T. Tocotrienol attenuates triglyceride accumulation in HepG2 cells and F344 rats. *Lipids.* 2012; 47(5): 471-81. [PMID: 22367056]

Takahashi A, Morita M, Yokoyama K, Suzuki T, Yamamoto T. Tob2 inhibits peroxisome proliferator-activated receptor γ2 expression by sequestering Smads and C/EBPα during adipocyte differentiation. *Mol Cell Biol.* 2012; 32(24): 5067-77. [PMID: 23071089]

2013

Ito M, Yamanashi Y, Toyoda Y, Izumi-Nakaseko H, Oda S,

- Sugiyama A, Kuroda M, Suzuki H, Takada T, Adachi-Akahane S. Disruption of Stard10 gene alters the PPAR α -mediated bile acid homeostasis. *Biochim Biophys Acta*. 2013; 1831(2):459-68. [PMID: 23200860]
- Yamashita-Sugahara Y, Tokuzawa Y, Nakachi Y, Kaneko-Yatsuka Y, Matsumoto M, Mizuno Y, Okazaki Y. Fam57b (family with sequence similarity 57, member B), a novel peroxisome proliferator-activated receptor γ target gene that regulates adipogenesis through ceramide synthesis. *J Biol Chem*. 2013; 288(7): 4522-37. [PMID: 23275342]
- Takada S, Kinugawa S, Hirabayashi K, Suga T, Yokota T, Takahashi M, Fukushima A, Homma T, Ono T, Sobrin MA, Masaki Y, Mizushima W, Kadoguchi T, Okita K, Tsutsui H. Angiotensin II receptor blocker improves the lowered exercise capacity and impaired mitochondrial function of the skeletal muscle in type 2 diabetic mice. *J Appl Physiol*. 2013; 114(7):844-57. [PMID: 23329824]
- Senmaru T, Fukui M, Okada H, Mineoka Y, Yamazaki M, Tsujikawa M, Hasegawa G, Kitawaki J, Obayashi H, Nakamura N. Testosterone deficiency induces markedly decreased serum triglycerides, increased small dense LDL, and hepatic steatosis mediated by dysregulation of lipid assembly and secretion in mice fed a high-fat diet. *Metabolism*. 2013 (Article in Press). [PMID: 23332447]
- Ichimura K, Kawashima Y, Nakamura T, Powell R, Hidoh Y, Terai S, Sakaida I, Kodera Y, Tsuji T, Ma JX, Sakai T, Matsumoto H, Obara T. Medaka fish, *Oryzias latipes*, as a model for human obesity-related glomerulopathy. *Biochem Biophys Res Commun*. 2013; 431(4):712-7. [PMID: 23353086]
- Fujii M, Shibasaki Y, Wakamatsu K, Honda Y, Kawauchi Y, Suzuki K, Arumugam S, Watanabe K, Ichida T, Asakura H, Yoneyama H. A murine model for non-alcoholic steatohepatitis showing evidence of association between diabetes and hepatocellular carcinoma. *Med Mol Morphol*. 2013 (Article in Press). [PMID: 23430399]
- Takata S, Sawa Y, Uchiyama T, Ishikawa H. Expression of Toll-Like Receptor 4 in Glomerular Endothelial Cells under Diabetic Conditions. *Acta Histochem Cytochem*. 2013; 46(1): 35-42. [PMID: 23554538]
- Arai S, Maehara N, Iwamura Y, Honda SI, Nakashima K, Kai T, Ogishi M, Morita K, Kurokawa J, Mori M, Motoi Y, Miyake K, Matsuhashi N, Yamamura KI, Ohara O, Shibuya A, Wakeland EK, Li QZ, Miyazaki T. Obesity-Associated Autoantibody Production Requires AIM to Retain the Immunoglobulin M Immune Complex on Follicular Dendritic Cells. *Cell Rep*. 2013; 3, 1-12. [PMID: 23562157]
- Iwata M, Ochi H, Hara Y, Tagawa M, Koga D, Okawa A, Asou Y. Initial Responses of Articular Tissues in a Murine High-Fat Diet-Induced Osteoarthritis Model: Pivotal Role of the IPFP as a Cytokine Fountain. *PLoS One*. 2013; 8(4): e60706. [PMID: 23593289]
- Yamagata H, Ikejima K, Takeda K, Aoyama T, Kon K, Okumura K, Watanabe S. Altered expression and function of hepatic natural killer T cells in obese and diabetic KK-A(y) mice. *Hepatol Res*. 2013 Mar;43(3):276-88. [PMID: 22834991]
- Aibara D, Matsusue K, Matsuo K, Takiguchi S, Gonzalez FJ, Yamano S. Expression of hepatic fat-specific protein 27 depends on the specific etiology of fatty liver. *Biol Pharm Bull*. 2013;36(11):1766-72.
- Aizawa N, Homma Y, Igawa Y. Influence of High Fat Diet Feeding for 20 Weeks on Lower Urinary Tract Function in Mice. *Low Urin Tract Symptoms*. 2013 May;5(2):101-8. [PMID: 26663378]
- Inoue H, Shintani N, Sakurai Y, Higashi S, Hayata-Takano A, Baba A, Hashimoto H. PACAP Inhibits β -cell Mass Expansion in a Mouse Model of Type II Diabetes: Persistent Suppressive Effects on Islet Density. *Front Endocrinol (Lausanne)*. 2013 Mar 11;4:27 [PMID: 23483824]
- Hashimoto T, Igarashi J, Hasan AU, Ohmori K, Kohno M, Nagai Y, Yamashita T, Kosaka H. Mifepristone promotes adiponectin production and improves insulin sensitivity in a mouse model of diet-induced obesity. *PLoS One*. 2013 Nov 6;8(11) [PMID: 24223187]
- Hasegawa K, Wakino S, Kimoto M, Minakuchi H, Fujimura K, Hosoya K, Komatsu M, Kaneko Y, Kanda T, Tokuyama H, Hayashi K, Itoh H. The hydrolase DDAH2 enhances pancreatic insulin secretion by transcriptional regulation of secretagogin through a Sirt1-dependent mechanism in mice. *FASEB J*. 2013 Jun;27(6):2301-15. [PMID: 23430976]
- Traber PG, Zomer E. Therapy of experimental NASH and fibrosis with galectin inhibitors. *PLoS One*. 2013 Dec 18;8(12):e83481 [PMID: 24367597]
- Uchiyama T, Takata S, Ishikawa H, Sawa Y. Altered dynamics in the renal lymphatic circulation of type 1 and type 2 diabetic mice. *Acta Histochem Cytochem*. 2013 Apr 30;46(2):97-104. [PMID: 23720608]
- Arai C, Miyake M, Matsumoto Y, Mizote A, Yoshizane C, Hanaya Y, Koide K, Yamada M, Hanaya T, Arai S, Fukuda S. Trehalose prevents adipocyte hypertrophy and mitigates insulin resistance in mice with established obesity. *J Nutr*

Sci Vitaminol (Tokyo). 2013;59(5):393-401. [PMID: 24418873]

Shimamura Y, Yoda M, Sakakibara H, Matsunaga K, Masuda S. Pu-erh tea suppresses diet-induced body fat accumulation in C57BL/6J mice by down-regulating SREBP-1c and related molecules. Biosci Biotechnol Biochem. 2013;77(7):1455-60. [PMID: 23832364]

Tanno S, Terao A, Okamatsu-Ogura Y, Kimura K. Hypothalamic prepro-orexin mRNA level is inversely correlated to the non-rapid eye movement sleep level in high-fat diet-induced obese mice. Obes Res Clin Pract. 2013 Jul-Aug;7(4):e251-7 [PMID: 24306152]

2014

Suga T, Kinugawa S, Takada S, Kadoguchi T, Fukushima A, Homma T, Masaki Y, Furihata T, Takahashi M, Sobrin MA, Ono T, Hirabayashi K, Yokota T, Tanaka S, Okita K, Tsutsui H. Combination of exercise training and diet restriction normalizes limited exercise capacity and impaired skeletal muscle function in diet-induced diabetic mice. Endocrinology. 2014 Jan;155(1):68-80. [PMID: 24189138]

Takakura K, Koido S, Fujii M, Hashiguchi T, Shibasaki Y, Yoneyama H, Katagi H, Kajihara M, Misawa T, Homma S, Ohkusa T, Tajiri H. Characterization of nonalcoholic steatohepatitis-derived hepatocellular carcinoma as a human stratification model in mice. Anticancer Res. 2014 Sep;34(9):4849-55. [PMID: 25202066]

Yamauchi A, Sakurai T, Kamiyoshi A, Ichikawa-Shindo Y, Kawate H, Igarashi K, Toriyama Y, Tanaka M, Liu T, Xian X, Imai A, Zhai L, Owa S, Arai T, Shindo T. Functional differentiation of RAMP2 and RAMP3 in their regulation of the vascular system. J Mol Cell Cardiol. 2014 Dec;77:73-85. [PMID: 25264174]

Ishikawa H, Takaki A, Tsuzaki R, Yasunaka T, Koike K, Shimomura Y, Seki H, Matsushita H, Miyake Y, Ikeda F, Shiraha H, Nouso K, Yamamoto K. L-carnitine prevents progression of non-alcoholic steatohepatitis in a mouse model with upregulation of mitochondrial pathway. PLoS One. 2014 Jul 1;9(7) [PMID: 24983359]

Kuramoto K, Sakai F, Yoshinori N, Nakamura TY, Wakabayashi S, Kojidani T, Haraguchi T, Hirose F, Osumi T. Deficiency of a lipid droplet protein, perilipin 5, suppresses myocardial lipid accumulation, thereby preventing type 1 diabetes-induced heart malfunction. Mol Cell Biol. 2014 Jul;34(14):2721-31. [PMID: 24820416]

Cheng J, Morisaki H, Toyama K, Sugimoto N, Shintani T, Tandelilin A, Hirase T, Holmes EW, Morisaki T. AMPD1: a

novel therapeutic target for reversing insulin resistance.

BMC Endocr Disord. 2014 Dec 15;14:96. [PMID: 25511531]

Naka S, Nomura R, Takashima Y, Okawa R, Ooshima T, Nakano K. A specific Streptococcus mutans strain aggravates non-alcoholic fatty liver disease. Oral Dis. 2014 Oct;20(7):700-6. [PMID: 25360469]

Takaki Y, Saito Y, Takasugi A, Toshimitsu K, Yamada S, Muramatsu T, Kimura M, Sugiyama K, Suzuki H, Arai E, Ojima H, Kanai Y, Saito H. Silencing of microRNA-122 is an early event during hepatocarcinogenesis from non-alcoholic steatohepatitis. Cancer Sci. 2014 Oct;105(10):1254-60. [PMID: 25117675]

Sawa Y, Takata S, Hatakeyama Y, Ishikawa H, Tsuruga E. Expression of toll-like receptor 2 in glomerular endothelial cells and promotion of diabetic nephropathy by Porphyromonas gingivalis lipopolysaccharide. PLoS One. 2014 May 16;9(5):e97165 [PMID: 24835775]

Yang W, Tanaka Y, Bundo M, Hirokawa N. Antioxidant signaling involving the microtubule motor KIF12 is an intracellular target of nutrition excess in beta cells. Dev Cell. 2014 Oct 27;31(2):202-14. [PMID: 25373778]

2015

Niwa M, Numaguchi Y, Ishii M, Kuwahata T, Kondo M, Shibata R, Miyata K, Oike Y, Murohara T. IRAP deficiency attenuates diet-induced obesity in mice through increased energy expenditure. Biochem Biophys Res Commun. 2015 Jan 30;457(1):12-8. [PMID: 25534853]

Nakano S, Katsuno K, Isaji M, Nagasawa T, Buehrer B, Walker S, Wilkison WO, Cheatham B. Remogliflozin Etabonate Improves Fatty Liver Disease in Diet-Induced Obese Male Mice. J Clin Exp Hepatol. 2015 Sep;5(3):190-8. [PMID: 26628836]

Tandelilin AA, Hirase T, Hudoyo AW, Cheng J, Toyama K, Morisaki H, Morisaki T. AMPD1 regulates mTORC1-p70 S6 kinase axis in the control of insulin sensitivity in skeletal muscle. BMC Endocr Disord. 2015;15:11. [PMID: 25887856]. *Erratum in: BMC Endocr Disord. 2015;8:15:80 [PMID: 26645205]

Nishimura W, Takahashi S, Yasuda K. MafA is critical for maintenance of the mature beta cell phenotype in mice. Diabetologia. 2015 Mar;58(3):566-74 [PMID: 25500951]

Konishi H, Shirabe K, Nakagawara H, Harimoto N, Yamashita Y, Ikegami T, Yoshizumi T, Soejima Y, Oda Y, Maehara Y. Suppression of silent information regulator 1 activity in noncancerous tissues of hepatocellular carcinoma: Possible association with non-B non-C hepatitis pathogenesis. Cancer Sci. 2015 May;106(5):542-9. [PMID: 25983359]

25736100]

Fushimi S, Myazawa F, Nakagawa K, Burdeos GC, Miyazawa T. Young Persimmon Ingestion Suppresses Lipid Oxidation in Rats. *J Nutr Sci Vitaminol (Tokyo)*. 2015;61(1):90-5. [PMID: 25994144]

Saito M, Kaneda A, Sugiyama T, Iida R, Otokuni K, Kaburagi M, Matsuoka H. Production of a mouse strain with impaired glucose tolerance by systemic heterozygous knockout of the glucokinase gene and its feasibility as a prediabetes model. *Exp Anim*. 2015;64(3):231-9. [PMID: 25765873]

Wang B, Tanaka K, Katsume T, Maruyama K, Ninomiya Y and Nenoi M. Dietary Modification of Mouse Response to Total-Body-Irradiation. *Evolution of Ionizing Radiation Research* September 17, 2015

Saito K, Uebano T, Maekawa K, Ishikawa M, Taguchi R, Nammo T, Nishimaki-Mogami T, Udagawa H, Fujii M, Shibasaki Y, Yoneyama H, Yasuda K, Saito Y. Characterization of hepatic lipid profiles in a mouse model with nonalcoholic steatohepatitis and subsequent fibrosis. *Sci Rep*. 2015 Aug 20;5:12466. [PMID: 26289793]

Takagi Y, Kinoshita K, Ozaki N, Seino Y, Murata Y, Oshida Y, Hayashi Y. Mice Deficient in Proglucagon-Derived Peptides Exhibit Glucose Intolerance on a High-Fat Diet but Are Resistant to Obesity. *PLoS One*. 2015 Sep 17;10(9):e0138322. [PMID: 26378455]

Sasaki T, Kinoshita Y, Matsui S, Kakuta S, Yokota-Hashimoto H, Kinoshita K, Iwasaki Y, Kinoshita T, Yada T, Amano N, Kitamura T. N-methyl-d-aspartate receptor coagonist d-serine suppresses intake of high-preference food. *Am J Physiol Regul Integr Comp Physiol*. 2015 Sep;309(5):R561-75. [PMID: 26157056]

Takahashi A, Adachi S, Morita M, Tokumasu M, Natsume T, Suzuki T, Yamamoto T. Post-transcriptional Stabilization of Ucp1 mRNA Protects Mice from Diet-Induced Obesity. *Cell Rep*. 2015 Dec 29;13(12):2756-67. [PMID: 26711342]

Uehara Y, Inoue M, Fukuda K, Yamakoshi H, Hosoi Y, Kanda H, Oshima M, Iwabuchi Y, Shibata H. Inhibition of β -catenin and STAT3 with a curcumin analog suppresses gastric carcinogenesis in vivo. *Gastric Cancer*. 2015 Oct;18(4):774-83. [PMID: 25331984]

Alixiding M, Aibibula Z, Iwata M, Piao J, Hara Y, Koga D, Okawa A, Morita S1, Asou Y. Pivotal role of Sirt6 in the crosstalk among ageing, metabolic syndrome and osteoarthritis. *Biochem Biophys Res Commun*. 2015 Oct 23;466(3):319-26 [PMID: 26362183]

Fujita T, Yamashita D, Uehara N, Inokuchi G, Hasegawa S, Otsuki N1, Nibu K. A high-fat diet delays age-related

hearing loss progression in C57BL/6J mice. *PLoS One*. 2015 Jan 27;10(1):e0117547 [PMID: 25625852]

2016

Higashimura Y, Naito Y, Takagi T, Uchiyama K, Mizushima K, Ushiroda C, Ohnogi H, Kudo Y, Yasui M, Inui S, Hisada T, Honda A, Matsuzaki Y, Yoshikawa T. Protective effect of agaric-oligosaccharides on gut dysbiosis and colon tumorigenesis in high-fat diet-fed mice. *Am J Physiol Gastrointest Liver Physiol*. 2016 Jan 14. [PMID: 26767984]

Noble EE, Kanoski SE. Early life exposure to obesogenic diets and learning and memory dysfunction. *Curr Opin Behav Sci*. 2016 Jun;9:7-14. [PMID: 26858972]

Yamato M, Kawano K, Yamanaka Y, Saiga M, Yamada K. TEMPOL increases NAD(+) and improves redox imbalance in obese mice. *Redox Biol*. 2016 Aug;8:316-22. [PMID: 26942863]

Takahashi T, Sugawara W, Takiguchi Y, Takizawa K, Nakabayashi A, Nakamura M, Nagano-Ito M, and Ichikawa S. Evidence-Based Complementary and Alternative Medicine Volume 2016 (2016), Article ID 4282758, 9. [PMID: None]

Kon K, Ikejima K, Morinaga M, Kusama H, Arai K, Aoyama T, Uchiyama A, Yamashina S, Watanabe S. L-carnitine prevents metabolic steatohepatitis in obese diabetic KK-Ay mice. *Hepatol Res*. 2016 Apr 7. [PMID: 27062266]

Hatanaka Y, Wada K, Kabuta T. Maternal high-fat diet leads to persistent synaptic instability in mouse offspring via oxidative stress during lactation. *Neurochem Int*. 2016 Jul;97:99-108. [PMID: 26970392]

Peñaa L, Meanaa, C, Astudilloa AM, Lordéna G, Valdearcosa M, Sato H, Murakamic M, Balsindea J, Balboaa MA. Critical role for cytosolic group IVA phospholipase A2 in early adipocyte differentiation and obesity. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids* Volume 1861, Issue 9, Part A, September 2016, Pages 1083–1095. [PMID: 27317983]

Kitada Y, Kajita K, Taguchi K, Mori I, Yamauchi M, Ikeda T, Kawashima M, Asano M, Kajita T, Ishizuka T, Banno Y, Kojima I, Chun J, Kamata S, Ishii I, Morita H. Blockade of Sphingosine 1-Phosphate Receptor 2 Signaling Attenuates High-Fat Diet-Induced Adipocyte Hypertrophy and Systemic Glucose Intolerance in Mice. *Endocrinology*. 2016 May;157(5):1839-51. [PMID: 26943364]

Takahashi K, Sasano T, Sugiyama K, Kurokawa J, Tamura N, Soejima Y, Sawabe M, Isobe M, Furukawa T. High-fat diet increases vulnerability to atrial arrhythmia by conduction disturbance via miR-27b. *J Mol Cell Cardiol*.

2016 Jan;90:38-46.[PMID: 26654778]

Xie G, Wang X, Liu P, Wei R, Chen W, Rajani C, Hernandez BY, Alegado R, Dong B, Li D, Jia W. Distinctly altered gut microbiota in the progression of liver disease. *Oncotarget*. 2016 Mar 29. [PMID: 27036035]

Sekine S, Yao A, Hattori K, Sugawara S, Naguro I, Koike M, Uchiyama Y, Takeda K, Ichijo H. The Ablation of Mitochondrial Protein Phosphatase Pgam5 Confers Resistance Against Metabolic Stress. *EBioMedicine*. 2016 Jan 29;5:82-92.[PMID: 27077115]

Aoki K, Tajima K, Taguri M, Terauchi Y. Effect of dehydroepiandrosterone (DHEA) on Akt and protein kinase C zeta (PKC ζ) phosphorylation in different tissues of C57BL6, insulin receptor substrate (IRS)1(-/-), and IRS2(-/-) male mice fed a high-fat diet. *J Steroid Biochem Mol Biol*. 2016 May;159:110-20.[PMID: 26976654]

Aibibula Z, Ailixiding M, Iwata M, Piao J, Hara Y, Okawa A, Asou Y. Xanthine oxidoreductase activation is implicated in the onset of metabolic arthritis. *Biochem Biophys Res Commun*. 2016 Mar 25;472(1):26-32.[PMID: 26903297]

Fukuda K, Uehara Y, Nakata E, Inoue M, Shimazu K, Yoshida T, Kanda H, Nanjo H, Hosoi Y, Yamakoshi H, Iwabuchi Y, Shibata H. A diarylpentanoid curcumin analog exhibits improved radioprotective potential in the intestinal mucosa. *Int J Radiat Biol*. 2016 Jul;92(7):388-94.[PMID: 27043482]

Shirai M, Arakawa S, Teranishi M, Kai K. Decreased hepatic phosphorylated p38 mitogen-activated protein kinase contributes to attenuation of thioacetamide-induced hepatic necrosis in diet-induced obese mice. *J Toxicol Sci*. 2016;41(2):245-53.[PMID: 26961609]

Kishida N, Matsuda S, Itano O, Shinoda M, Kitago M, Yagi H, Abe Y, Hibi T, Masugi Y, Aiura K, Sakamoto M, Kitagawa Y. Development of a novel mouse model of hepatocellular carcinoma with nonalcoholic steatohepatitis using a high-fat, choline-deficient diet and intraperitoneal injection of diethylnitrosamine. *BMC Gastroenterol*. 2016 Jun 13;16(1):61.[PMID: 27296438]

Nakazono S, Cho K, Isaka S, Abu R, Yokose T, Murata M, Ueno M, Tachibana K, Hirasaka K, Kim D, Oda T. Anti-obesity effects of enzymatically-digested alginate oligomer in mice model fed a high-fat-diet. *Bioactive Carbohydrates and Dietary Fibre* Vol7,2, April 2016, 1-8 [PMID: none]

Takahashi T, Sugawara W, Takiguchi Y, Takizawa K, Nakabayashi A, Nakamura M, Nagano-Ito M, Ichikawa S. Identification of Plants That Inhibit Lipid Droplet Formation in Liver Cells: Rubus suavissimus Leaf Extract

Protects Mice from High-Fat Diet-Induced Fatty Liver by Directly Affecting Liver Cells. *Evid Based Complement Alternat Med*. 2016;2016:4282758.[PMID: 27429636]

Toita R, Kawano T, Murata M, Kang JH. Anti-obesity and anti-inflammatory effects of macrophage-targeted interleukin-10-conjugated liposomes in obese mice. *Biomaterials*. 2016 Dec;110:81-88.[PMID: 27716475]

Yamamoto S, Oshima Y, Saitou T, Watanabe T, Miyake T, Yoshida O, Tokumoto Y, Abe M, Matsuura B, Hiasa Y, Imamura T. Quantitative imaging of fibrotic and morphological changes in liver of non-alcoholic steatohepatitis (NASH) model mice by second harmonic generation (SHG) and auto-fluorescence (AF) imaging using two-photon excitation microscopy (TPEM) *Biochemistry and Biophysics Reports* 8 (2016) 277–283 [PMID: none]

Nagaishi K, Mizue Y, Chikenji T, Otani M, Nakano M, Konari N, Fujimiya M. Mesenchymal stem cell therapy ameliorates diabetic nephropathy via the paracrine effect of renal trophic factors including exosomes. *Sci Rep*. 2016 Oct 10:6:34842.[PMID: 27721418]

Ijuin T, Hosooka T, Takenawa T. Phosphatidylinositol 3,4,5-Trisphosphate Phosphatase SKIP Links Endoplasmic Reticulum Stress in Skeletal Muscle to Insulin Resistance. *Mol Cell Biol*. 2015 Oct 19;36(1):108-18.[PMID: 26483413]

Akiko Takashima, Kimihiko Sano, Shiho Nakagawa, Gen Toshima, Junichiro Takahashi, Fumika Kurosaki, Akira Sasaki, Kazuyuki Hiwatashi, Keishi Hata. In vitro and in vivo antidiabetic effects of the ethanol extract from Lentinula edodes (Shiitake). *Nutrafoods* (2016) 15:279-284 [PMID: None]

2017

Kodo K, Sugimoto S, Nakajima H, Mori J, Itoh I, Fukuhara S, Shigehara K, Nishikawa T, Kosaka K, Hosoi H. Erythropoietin (EPO) ameliorates obesity and glucose homeostasis by promoting thermogenesis and endocrine function of classical brown adipose tissue (BAT) in diet-induced obese mice. *PLoS One*. 2017 Mar 13;12(3):e0173661 [PMID: 28288167]

Tomiga Y, Yoshimura S, Ito A, Nakashima S, Kawanaka K, Uehara Y, Tanaka H, Higaki Y. Exercise training rescues high fat diet-induced neuronal nitric oxide synthase expression in the hippocampus and cerebral cortex of mice. *Nitric Oxide*. 2017 Jun 1;66:71-77.[PMID: 28302517]

Kon K, Ikejima K, Morinaga M, Kusama H, Arai K, Aoyama T, Uchiyama A, Yamashina S, Watanabe S. L-carnitine prevents metabolic steatohepatitis in obese diabetic KK-Ay

- mice. *Hepatol Res.* 2017 Mar;47(3):E44-E54. [PMID: 27062266]
- Yamamoto R, Iida A, Tanikawa K, Shiratsuchi H, Tokuda M, Matsui T, Nakamura T. Dietary D-Allose Ameliorates Hepatic Inflammation in Mice with Non-alcoholic Steatohepatitis. *Food Science and Technology Research* Vol. 23 (2017) No. 2 p. 319-327 [PMID: None]
- Hino J, Nakatani M, Arai Y, Tsuchida K, Shirai M, Miyazato M, Kangawa K. Overexpression of bone morphogenetic protein-3b (BMP-3b) in adipose tissues protects against high-fat diet-induced obesity. *Int J Obes (Lond)*. 2017 Apr;41(4):483-488.
- Afrin MR, Arumugam S, Rahman MA, Karuppagounder V, Harima M, Suzuki H, Miyashita S, Suzuki K, Ueno K, Yoneyama H, Watanabe K. Curcumin reduces the risk of chronic kidney damage in mice with nonalcoholic steatohepatitis by modulating endoplasmic reticulum stress and MAPK signaling. *Int Immunopharmacol*. 2017 Jun 5;49:161-167.[PMID: 28595079]
- Ohuchida K, Sada M, Abe T, Endo S, Koikawa K, Iwamoto C, Miura D, Mizuuchi Y, Moriyama T, Nakata K, Miyasaka Y, Manabe T, Ohtsuka T, Nagai E, Mizumoto K, Oda Y, Hashizume M, Nakamura M. Extra-pancreatic invasion induces lipolytic and fibrotic changes in the adipose microenvironment, with released fatty acids enhancing the invasiveness of pancreatic cancer cells. *Oncotarget*. 2017 Mar 14;8(11):18280-18295.[PMID: 28407685]
- Takemoto M, Takemoto H, Kouno H, Soeda T, Moriya T. A Simple, Enzymatic Biotransformation Method Using Fresh Green Tea Leaves Efficiently Generates Theaflavin-Containing Fermentation Water That Has Potent Physiological Functions in Mice and Humans. *Biol Pharm Bull*. 2017 Jun 1;40(6):860-866.[PMID: 28381812]
- Sreedhar R, Arumugam S, Thandavarayan RA, Karuppagounder V, Koga Y, Nakamura T, Harima M, Watanabe K. Role of 14-3-3 η protein on cardiac fatty acid metabolism and macrophage polarization after high fat diet induced type 2 diabetes mellitus.[PMID: 28483670]
- Kubo H, Hoshi M, Matsumoto T, Irie M, Oura S, Tsutsumi H, Hata Y, Yamamoto Y, Saito K. Sake lees extract improves hepatic lipid accumulation in high fat diet-fed mice. *Lipids Health Dis*. 2017 Jun 3;16(1):106. [PMID: 28578672]
- Lye HS, Kato T, Low WY, Taylor TD, Prakash T, Lew LC, Ohno H, Liong MT. Lactobacillus fermentum FTDC 8312 combats hypercholesterolemia via alteration of gut microbiota. *J Biotechnol*. 2017 Nov 20;262:75-83. [PMID: 28935567]
- Bae CR, Hino J, Hosoda H, Arai Y, Son C, Makino H, Tokudome T, Tomita T, Kimura T, Nojiri T, Hosoda K, Miyazato M, Kangawa K. Overexpression of C-type Natriuretic Peptide in Endothelial Cells Protects against Insulin Resistance and Inflammation during Diet-induced Obesity. *Sci Rep*. 2017 Aug 29;7(1):9807.[PMID: 28852070]
- Kusumoto Y, Irie J, Iwabu K, Tagawa H, Itoh A, Kato M, Kobayashi N, Tanaka K, Kikuchi R, Fujita M, Nakajima Y, Morimoto K, Sugizaki T, Yamada S, Kawai T, Watanabe M, Oike Y, Itoh H. Bile acid binding resin prevents fat accumulation through intestinal microbiota in high-fat diet-induced obesity in mice. *Metabolism*. 2017 Jun;71:1-6.[PMID: 28521862]
- Komori T, Morikawa Y. Oncostatin M in the development of metabolic syndrome and its potential as a novel therapeutic target. *Anat Sci Int*. 2017 Nov 4.[PMID: 29103176]
- Saito T, Muramatsu M, Ishii Y, Saigo Y, Konuma T, Toriniwa Y, Miyajima K, Ohta T. Pathophysiological analysis of the progression of hepatic lesions in STAM mice. *Physiol Res*. 2017 Nov 24;66(5):791-799.[PMID: 28730823]
- Sasaki T, Yasoshima Y, Matsui S, Yokota-Hashimoto H, Kobayashi M, Kitamura T. Intraperitoneal injection of d-serine inhibits high-fat diet intake and preference in male mice. *Appetite*. 2017 Nov 1;118:120-128.[PMID: 28803978]
- Hirose S, Asano K, Nakane A. Attenuation of obesity-induced inflammation in mice orally administered with salmon cartilage proteoglycan, a prophylactic agent. *Biochem Biophys Res Commun*. 2017 Mar 11;484(3):480-485. [PMID: 28089867]
- Hirotani Y, Fukamachi J, Ueyama R, Urashima Y, Ikeda K. Effects of Capsaicin Coadministered with Eicosapentaenoic Acid on Obesity-Related Dysregulation in High-Fat-Fed Mice. *Biol Pharm Bull*. 2017;40(9):1581-1585.[PMID: 28867743]
- Pogribny IP, Dreval K, Kindrat I, Melnyk S, Jimenez L, de Conti A, Tryndyak V, Pogribna M, Ortega JF, James SJ, Rusyn I, Beland FA. Epigenetically mediated inhibition of S-adenosylhomocysteine hydrolase and the associated dysregulation of 1-carbon metabolism in nonalcoholic steatohepatitis and hepatocellular carcinoma. *FASEB J*. 2017 Nov 10. pii: fj.201700866R.[PMID: 29127188]
- Afrin R, Arumugam S, Rahman A, Wahed MI, Karuppagounder V, Harima M, Suzuki H, Miyashita S, Suzuki K, Yoneyama H, Ueno K, Watanabe K. Curcumin ameliorates liver damage and progression of NASH in NASH-HCC mouse model possibly by modulating HMGB1-NF- κ B translocation. *Int Immunopharmacol*.

2017 Mar;44:174-182. [PMID: 28110063]

Teranishi H, Hayashi M, Higa R, Mori K, Miyazawa T, Hino J, Amano Y, Tozawa R, Ida T, Hanada T, Miyazato M, Hanada R, Kangawa K, Nakao K. Role of neuromedin U in accelerating of non-alcoholic steatohepatitis in mice. *Peptides.* 2017 Oct 7. pii: S0196-9781(17)30294-2. [PMID: 29017855]

Takahashi Y, Matsunaga Y, Tamura Y, Terada S, Hatta H. Pre-Exercise High-Fat Diet for 3 Days Affects Post-Exercise Skeletal Muscle Glycogen Repletion. *J Nutr Sci Vitaminol (Tokyo).* 2017;63(5):323-330.[PMID: 29225317]

Suda M, Shimizu I, Yoshida Y, Hayashi Y, Ikegami R, Katsuumi G, Wakasugi T, Yoshida Y, Okuda S, Soga T, Minamino T. Inhibition of dipeptidyl peptidase-4 ameliorates cardiac ischemia and systolic dysfunction by up-regulating the FGF-2/EGR-1 pathway. *PLoS One.* 2017 Aug 3;12(8):e0182422.[PMID: 28771625]

de Conti A, Ortega JF, Tryndyak V, Dreval K, Moreno FS, Rusyn I, Beland FA, Pogribny IP. MicroRNA deregulation in nonalcoholic steatohepatitis-associated liver carcinogenesis. *Oncotarget.* 2017 Aug 1;8(51):88517-88528. [PMID: 29179453]

Ideta T, Shirakami Y, Ohnishi M, Maruta A, Obara K, Miyazaki T, Kochi T, Sakai H, Tomita H, Tanaka T, Blaner WS, Shimizu M. Non-alcoholic steatohepatitis-related liver tumorigenesis is suppressed in mice lacking hepatic retinoid storage. *Oncotarget.* 2017 Aug 7;8(41):70695-70706.[PMID: 29050312]

Sugizaki T, Zhu S, Guo G, Matsumoto A, Zhao J, Endo M, Horiguchi H, Morinaga J, Tian Z, Kadomatsu T, Miyata K, Itoh H, Oike Y. Treatment of diabetic mice with the SGLT2 inhibitor TA-1887 antagonizes diabetic cachexia and decreases mortality. *NPJ Aging Mech Dis.* 2017 Sep 8;3:12.[PMID: 28900540]

Takemoto M, Takemoto H, Kouno H, Soeda T, Moriya T. A Simple, Enzymatic Biotransformation Method Using Fresh Green Tea Leaves Efficiently Generates Theaflavin-Containing Fermentation Water That Has Potent Physiological Functions in Mice and Humans. *Biol Pharm Bull.* 2017 Jun 1;40(6):860-866.[PMID: 28381812]

Shimizu T, Sato T, Tsukiyama K, Fujita H, Kato S, Hoizumi M, Shirasawa H, Narita T, Terada Y, Seino Y3, Yamada Y. Food Intake Affects Sperm-Egg Fusion Through the GIP/PSG17 Axis in Mice. *Endocrinology.* 2017 Jul 1;158(7):2134-2144.[PMID: 28430907]

2018

Satoh M, Iwabuchi K. Role of Natural Killer T Cells in the Development of Obesity and Insulin Resistance: Insights From Recent Progress. *Front Immunol.* 2018 Jun 11:9:1314.[PMID: 29942311]

Higashi Y, Yamakuchi M, Fukushige T, Ibusuki A, Hashiguchi T, Kanekura T. High-fat diet exacerbates imiquimod-induced psoriasis-like dermatitis in mice. *Exp Dermatol.* 2018 Feb;27(2):178-184. [PMID: 29247486]

Al-Fayoumi S, Hashiguchi T, Shirakata Y, Mascarenhas J, Singer JW. Pilot study of the antifibrotic effects of the multikinase inhibitor pacritinib in a mouse model of liver fibrosis. *J Exp Pharmacol.* 2018 May 9;10:9-17.[PMID: 29785143]

Yoshida Y, Furukawa JI, Naito S, Higashino K, Numata Y, Shinohara Y. Identification of unique glycoisoforms of vitamin D-binding protein and haptoglobin as biomarker candidates in hepatocarcinogenesis of STAM mice. *Glycoconj J.* 2018 Sep 8.[PMID: 30194503]

Maejima Y, Yokota S, O'Hashi R, Aoyama M, Shimomura K. The effect of avertin anesthesia and a mixture of three types of anesthetic agents on food intake and body weight in high fat-induced obese male and female mice. *Exp Anim.* 2018 Sep 13.[PMID: 30210091]

Lo Re O, Panebianco C, Porto S, Cervi C, Rappa F, Di Biase S, Caraglia M, Pazienza V, Vinciguerra M. Fasting inhibits hepatic stellate cells activation and potentiates anti-cancer activity of Sorafenib in hepatocellular cancer cells. *J Cell Physiol.* 2018 Feb;233(2):1202-1212.[PMID: 28471474]

Fujisawa K, Takami T, Fukui Y, Nagatomo T, Saeki I, Matsumoto T, Hidaka I, Yamamoto N, Okamoto T, Furutani-Seiki M, Sakaida I. Assessment of high fat diet-induced fatty liver in medaka. *Biol Open.* 2018 Aug 20.[PMID: 30127096]

Akaki J, Tachi S, Nakamura N, Arai T, Yamasaki H, Inoue M, Makino T. Promotive effect of Bofutsushosan (Fangfengtongshengsan) on lipid and cholesterol excretion in feces in mice treated with a high-fat diet. *J Ethnopharmacol.* 2018 Jun 28;220:1-8.[PMID: 29588201]

Sasaki T, Numano R, Yokota-Hashimoto H, Matsui S, Kimura N, Takeuchi H, Kitamura T. A central-acting connexin inhibitor, INI-0602, prevents high-fat diet-induced feeding pattern disturbances and obesity in mice. *Mol Brain.* 2018 May 24;11(1):28.[PMID: 29793524]

Mori T, Suzuki-Yamazaki N, Takaki S. Lnk/Sh2b3 Regulates Adipose Inflammation and Glucose Tolerance through Group 1 ILCs. *Cell Rep.* 2018 Aug 14;24(7):1830-1841.



日本クレア株式会社

東京 A D 部 TEL. 03-5704-7050 FAX. 03-3792-2032
大阪 A D 部 TEL. 06-4861-7101 FAX. 06-4861-7108

<http://www.CLEA-Japan.com>

仙台出張所 TEL. 022-352-4417 FAX. 022-352-4419
札幌出張所 TEL. 011-631-2725 FAX. 011-644-9209

[PMID: 30110639]

Oishi K, Ohyama S, Higo-Yamamoto S. Chronic sleep disorder induced by psychophysiological stress induces glucose intolerance without adipose inflammation in mice.

Biochem Biophys Res Commun. 2018 Jan 22;495(4):2616-2621. [PMID: 29288667]

Homeyer A, Hammad S, Schwen LO, Dahmen U, Höfener H, Gao Y, Dooley S, Schenk A. Focused scores enable reliable discrimination of small differences in steatosis. Diagn Pathol. 2018 Sep 20;13(1):76. [PMID: 30231920]

Yoshida K, Kita Y, Tokuoka SM, Hamano F, Yamazaki M, Sakimura K, Kano M, Shimizu T. Monoacylglycerol lipase deficiency affects diet-induced obesity, fat absorption, and feeding behavior in CB1 cannabinoid receptor-deficient mice. FASEB J. 2018 Sep 28:fj201801203R. [PMID: 30265576]

Nakashima A, Sugimoto R, Suzuki K, Shirakata Y, Hashiguchi T, Yoshida C, Nakano Y. Anti-fibrotic activity of Euglena gracilis and paramylon in a mouse model of non-alcoholic steatohepatitis. Food Sci Nutr. 2018 Nov 8;7(1):139-147. [PMID: 30680167]

Ardiansyah, Inagawa Y, Koseki T, Agista AZ, Ikeda I, Goto T, Komai M, Shirakawa H. Adenosine and adenosine-5'-monophosphate ingestion ameliorates abnormal glucose metabolism in mice fed a high-fat diet. BMC Complement Altern Med. 2018 Nov 14;18(1):304. [PMID: 30428888]

Naka S, Wato K, Hatakeyama R, Okawa R, Nomura R, Nakano K. Longitudinal comparison of Streptococcus mutans-induced aggravation of non-alcoholic steatohepatitis in mice. J Oral Microbiol. 2018 Jan 22;10(1):1428005. [PMID: 29503703]

Dechassa ML, Tryndyak V, de Conti A, Xiao W, Beland FA, Pogribny IP. Identification of chromatin-accessible domains in non-alcoholic steatohepatitis-derived hepatocellular carcinoma. Mol Carcinog. 2018 Aug;57(8):978-987. [PMID: 29603380]

Kawaguchi T, Nakano D, Koga H, Torimura T. Effects of a DPP4 Inhibitor on Progression of NASH-related HCC and the p62/ Keap1/Nrf2-Pentose Phosphate Pathway in a Mouse Model. Liver Cancer. 2018, DOI_10.1159_000491763 [PMID: None]

Nishina S, Yamauchi A, Kawaguchi T, Kaku K, Goto M, Sasaki K, Hara Y, Tomiyama Y, Kuribayashi F, Torimura T, Hino K. Dipeptidyl Peptidase 4 Inhibitors Reduce Hepatocellular Carcinoma by Activating Lymphocyte Chemotaxis in Mice. Cell Mol Gastroenterol Hepatol. 2018 Sep 11;7(1):115-134. [PMID: 30510994]

Wakabayashi C, Kunugi H. Involvement of IL-6 and GSK3β in impaired sensorimotor gating induced by high-fat diet. Neurosci Res. 2018 Oct 13. pii: S0168-0102(18)30328-6. [PMID: 30326250]

Yoshida T, Maruyama T, Miura M, Inoue M, Fukuda K, Shimazu K, Taguchi D, Kanda H, Oshima M, Iwabuchi Y, Shibata H. Dietary intake of pyrolyzed deketene curcumin inhibits gastric carcinogenesis. Journal of Functional Foods Volume 50, November 2018, Pages 192-200 [PMID: None]

Liu T, Kamiyoshi A, Tanaka M, Iida S, Sakurai T, Ichikawa-Shindo Y, Kawate H, Hirabayashi K, Dai K, Cui N, Tanaka M, Wei Y, Nakamura K, Matsui S, Yamauchi A, Shindo T. RAMP3 deficiency enhances postmenopausal obesity and metabolic disorders. Peptides. 2018 Oct 29;110:10-18. [PMID: 30487357]

2019

Komatsu G, Nonomura T, Sasaki M, Ishida Y, Arai S, Miyazaki T. AIM-deficient mouse fed a high-trans fat, high-cholesterol diet: a new animal model for nonalcoholic fatty liver disease. Exp Anim. 2019; 68(2): 147-158. [PMID: 30487357]

Tsuma Y, Mori J, Ota T, Kawabe Y, Morimoto H, Fukuhara S, Kodo K, Umemura A, Nakajima H, Hosoi H. Erythropoietin and long-acting erythropoiesis stimulating agent ameliorate non-alcoholic fatty liver disease by increasing lipolysis and decreasing lipogenesis via EPOR/STAT pathway. Biochem Biophys Res Commun. 2019 Jan 29;509(1):306-313. [PMID: 30583863]

Murakami-Nishida S, Matsumura T, Senokuchi T, Ishii N, Kinoshita H, Yamada S, Morita Y, Nishida S, Motoshima H, Kondo T, Komohara Y, Araki E. Pioglitazone suppresses macrophage proliferation in apolipoprotein-E deficient mice by activating PPARγ. Atherosclerosis. 2019 Jul;286:30-39. [PMID: 31096071]

Borowa-Mazgaj B, de Conti A, Tryndyak V, Steward CR, Jimenez L, Melnyk S, Seneshaw M, Mirshahi F, Rusyn I, Beland FA, Sanyal AJ, Pogribny IP. Gene expression and DNA methylation alterations in the glycine N-methyltransferase gene in diet-induced nonalcoholic fatty liver disease-associated carcinogenesis. Toxicol Sci. 2019 May 14. [PMID: 31086990]

Yoshida Y, Shimizu I, Hayashi Y, Ikegami R, Suda M, Katsumi G, Wakasugi T, Nakao M, Nakagami H, Morishita R, Minamino T. Peptide vaccine for semaphorin3E ameliorates systemic glucose intolerance in mice with dietary obesity. Sci Rep. 2019 Mar 7;9(1):3858.

[PMID: 30846754]

Hye Khan MA, Schmidt J, Stavniichuk A, Imig JD, Merk D.

A dual farnesoid X receptor/soluble epoxide hydrolase modulator treats non-alcoholic steatohepatitis in mice. Biochem Pharmacol. 2019 Aug;166:212-221. [PMID: 31129048]

Kawakami T, Takasaki S, Kadota Y, Fukuoka D, Sato M, Suzuki S. Regulatory role of metallothionein-1/2 on development of sex differences in a high-fat diet-induced obesity. Life Sci. 2019 Jun 1;226:12-21. [PMID: 30954474]

Ushiroda C, Naito Y, Takagi T, Uchiyama K, Mizushima K, Higashimura Y, Yasukawa Z, Okubo T, Inoue R, Honda A, Matsuzaki Y, Itoh Y, Green tea polyphenol (epigallocatechin-3-gallate) improves gut dysbiosis and serum bile acids dysregulation in high-fat diet-fed mice. J Clin Biochem Nutr. 2019 Apr, 1-13. [PMID: None]

Shimoda H, Tanaka J, Zaiki K, Nakaoji K Tiliroside, a Constituent of Strawberry Seeds, Slightly Enhances Hyaluronan Production via Hyaluronan Synthase 2 Expression in Mouse Skin and Human Fibroblasts. Int J of Biomedical Science 15(1), 24-31, Mar 15, 2019 [PMID: None]

Matsumoto K, Shimada H, Morishima H, Akahori R, Kida N, Resistant Starch - Supplemented Udon Noodles Prevent Impaired Glucose Tolerance and Induce Intestinal Immunoglobulin - A Secretion in Mice Starch 05 June 2019. [PMID: None]

Iida A, Kuranuki S, Yamamoto R, Uchida M, Ohta M, Ichimura M, Tsuneyama K, Masaki T, Seike M, Nakamura T. Analysis of amino acid profiles of blood over time and biomarkers associated with non-alcoholic steatohepatitis in STAM mice. Exp Anim. 2019 Jun 1. [PMID: 31155606]

Zhu S, Tian Z, Torigoe D, Zhao J, Xie P, Sugizaki T, Sato M, Horiguchi H, Terada K, Kadomatsu T, Miyata K, Oike Y. Aging- and obesity-related peri-muscular adipose tissue accelerates muscle atrophy. PLoS One. 2019 Aug 23;14(8):e0221366. [PMID: 31442231]

Shikama Y, Kurosawa M, Furukawa M, Ishimaru N, Matsushita K. Involvement of adiponectin in age-related increases in tear production in mice. Aging (Albany NY). 2019 Oct 8;11(19):8329-8346. [PMID: 31596727]

Kawaguchi T, Nakano D, Koga H, Torimura T. Effects of a DPP4 Inhibitor on Progression of NASH-related HCC and the p62/ Keap1/Nrf2-Pentose Phosphate Pathway in a Mouse Model. Liver Cancer. 2019 Oct;8(5):359-372. [PMID: 31768345]

2020

Kenichi Watanabe, Mst. Rejina Afrin, Remya Sreedhar, Vengadeshprabhu Karuppagounder, Meilei Harima, Xavier Alexander, Ravichandiran Velayutham, Somasundaram Arumugam. Pharmacological Investigation of Ceraceomyces tessulatus (Agaricomycetes) in Mice with Non-Alcoholic Steatohepatitis. [Forth coming] Shuhei Nishida, Takeshi Matsumura, Takafumi Senokuchi, Saiko Murakami-Nishida, Norio Ishii, Yutaro Morita, Yoshitaka Yagi, Hiroyuki Motoshima, Tatsuya Kondo, Eiichi Araki. Inhibition of inflammation-mediated DPP-4 expression by linagliptin increases M2 macrophages in atherosclerotic lesions. Biochem Biophys Res Commun. 2020 Mar 26;524(1):8-15. [PMID: 31964532]

Toshiaki Teratani, Kengo Tomita, Sachiko Toma-Fukai, Yutaro Nakamura, Toshimasa Itoh, Hikaru Shimizu, Yasunaga Shiraishi, Nao Sugihara, Masaaki Higashiyama, Takahiko Shimizu, Ikuo Inoue, Yasuhiro Takenaka, Ryota Hokari, Takeshi Adachi, Toshiyuki Shimizu, Soichiro Miura, Takanori Kanai. Redox-dependent PPAR γ /Tnpo1 complex formation enhances PPAR γ nuclear localization and signaling. Free Radic Biol Med. 2020 Jun 15;156:45-56. [PMID: 32553752]

Yusuke Sasaki, Masato Asahiyama, Toshiya Tanaka, Shogo Yamamoto, Kentaro Murakami, Wakana Kamiya, Yoshihiro Matsumura, Tsuyoshi Osawa, Motonobu Anai, Jean-Charles Fruchart, Hiroyuki Aburatani, Juro Sakai, Tatsuhiko Kodama. Pemafibrate, a selective PPAR α modulator, prevents non-alcoholic steatohepatitis development without reducing the hepatic triglyceride content. Sci Rep. 2020 May 8;10(1):7818. [PMID: 32385406]

Masahiro Matsui, Shinya Fukunishi, Takashi Nakano, Takaaki Ueno, Kazuhide Higuchi, Akira Asai. Ileal bile acid transporter inhibitor improves hepatic steatosis by changing the gut microbiota dysbiosis in NAFLD model mice. <Research Square preprint.> [PMID: none]

Yuko Maejima, Shoko Yokota, Shoichiro Horita, Kenju Shimomura. Early life high-fat diet exposure evokes normal weight obesity. Nutr Metab (Lond). 2020 Jun 24;17:48. [PMID: 32595758]

Shohei Tanaka, Yasuhiro Nemoto, Yuria Takei, Ryo Morikawa, Shigeru Oshima, Takashi Nagaishi, Ryuichi Okamoto, Kiichiro Tsuchiya, Tetsuya Nakamura, Susanne Stutte, Mamoru Watanabe. High-fat diet-derived free fatty acids impair the intestinal immune system and increase sensitivity to intestinal epithelial damage. Biochem

- Biophys Res Commun. 2020 Feb 19;522(4):971-977. [PMID: 31810607]
- Hiroyasu Sato, Yoshitaka Taketomi, Yoshimi Miki, Remi Murase, Kei Yamamoto, Makoto Murakami. Secreted Phospholipase PLA2G2D Contributes to Metabolic Health by Mobilizing ω 3 Polyunsaturated Fatty Acids in WAT. Cell Rep. 2020 May 5;31(5):107579. [PMID: 32375030]
- Takeshi Moriishi, Ryosuke Ozasa, Takuya Ishimoto, Takayoshi Nakano, Tomoka Hasegawa, Toshihiro Miyazaki, Wenguang Liu, Ryo Fukuyama, Yuying Wang, Hisato Komori, Xin Qin, Norio Amizuka, Toshihisa Komori. Osteocalcin is necessary for the alignment of apatite crystallites, but not glucose metabolism, testosterone synthesis, or muscle mass. PLoS Genet. 2020 May; 16(5): e1008586. [PMID: 32463816]
- Momoka Nishimura, Yuki Nomura, Moritoki Egi, Norihiko Obata, Makoto Tsunoda, Satoshi Mizobuchi. Suppression of behavioral activity and hippocampal noradrenaline caused by surgical stress in type 2 diabetes model mice. BMC Neurosci. 2020 Feb 17;21(1):8. [PMID: 32066381]
- Chikako Arai, Aki Suyama, Shigeyuki Arai, Norie Arai, Chiyo Yoshizane, Satomi Koya-Miyata, Akiko Mizote, Shin Endo, Toshio Ariyasu, Hitoshi Mitsuzumi, Shimpei Ushio. Trehalose itself plays a critical role on lipid metabolism: Trehalose increases jejunum cytoplasmic lipid droplets which negatively correlated with mesenteric adipocyte size in both HFD-fed trehalase KO and WT mice. Nutr Metab (Lond). 2020 Mar 18;17:22. [PMID: 32206077]
- Shin Sato, Toshio Norikura, Yuuka Mukai, Shin Yamaoka, Keigo Mikame. Lignin-derived low-molecular-weight oxidized lignophenol stimulates AMP-activated protein kinase and suppresses renal inflammation and interstitial fibrosis in high fat diet-fed mice. Chem Biol Interact. 2020 Feb 25;318:108977. [PMID: 32035863]
- Tomoaki Inazumi, Kiyotaka Yamada, Naritoshi Shirata, Hiroyasu Sato, Yoshitaka Taketomi, Kazunori Morita, Hirofumi Hohjoh, Soken Tsuchiya, Kentaro Oniki, Takehisa Watanabe, Yutaka Sasaki, Yuichi Oike, Yasuhiro Ogata, Junji Saruwatari, Makoto Murakami, Yukihiko Sugimoto. Prostaglandin E 2-EP4 Axis Promotes Lipolysis and Fibrosis in Adipose Tissue Leading to Ectopic Fat Deposition and Insulin Resistance. Cell Rep. 2020 Oct 13;33(2):108265. [PMID: 33053354]
- Eri Nanizawa, Yuki Tamaki, Reika Sono, Rintaro Miyashita, Yumi Hayashi, Ayumu Kanbe, Hiroyasu Ito, Tetsuya Ishikawa. Short-term high-fat diet intake leads to exacerbation of concanavalin A-induced liver injury through the induction of procoagulation state. Biochem Biophys Rep 2020 Feb 12;22:100736. [PMID: 32083190]
- Kazuya Okushin, Takeya Tsutsumi, Kazuhiko Ikeuchi, Akira Kado, Kenichiro Enooku, Hidetaka Fujinaga, Naoko Yamauchi, Tetsuo Ushiku, Kyoji Moriya, Hiroshi Yotsuyanagi, Kazuhiko Koike. Heterozygous knockout of Bile salt export pump ameliorates liver steatosis in mice fed a high-fat diet. PLoS One. 2020 Aug 12;15(8):e0234750. [PMID: 32785220]
- Yuka Sasaki, Hiroshi Kuwata, Moe Akatsu, Yuri Yamakawa, Tsubasa Ochiai, Emiko Yoda, Yoshihito Nakatani, Chieko Yokoyama, Shuntaro Hara. Involvement of prostacyclin synthase in high-fat-diet-induced obesity. Prostaglandins Other Lipid Mediat. 2020 Dec 28;153:106523. [PMID: 33383181]
- Masahiro Matsui, Shinya Fukunishi, Takashi Nakano, Takaaki Ueno, Kazuhide Higuchi, Akira Asai. Ileal bile acid transporter inhibitor improves hepatic steatosis by changing the gut microbiota dysbiosis in NAFLD model mice. Research Square (Preprint) [PMID: none]
- Koichi Sugimoto, Tetsuro Yokokawa, Tomofumi Misaka, Takashi Kaneshiro, Akiomi Yoshihisa, Kazuhiko Nakazato, Yasuchika Takeishi. High-Fat Diet Attenuates the Improvement of Hypoxia-induced Pulmonary Artery Hypertension in Mice During Reoxygenation. Research Square (Preprint) [PMID: none]
- Norihiro Nagai, Hirohiko Kawashima, Eriko Toda, Kohei Homma, Hideto Osada, Naymel A Guzman, Shinsuke Shibata, Yasuo Uchiyama, Hideyuki Okano, Kazuo Tsubota, Yoko Ozawa. Renin-angiotensin system impairs macrophage lipid metabolism to promote age-related macular degeneration in mouse models. Commun Biol. 2020 Dec 9;3(1):767. [PMID: 33299105]
- Kenichi Watanabe, Rejina Afrin, Remya Sreedhar, Vengadeshprabhu Karuppagounder, Meilei Harima, Xavier Alexander, Ravichandiran Velayutham, Somasundaram Arumugam. Pharmacological Investigation of Ceraceomyces tessulatus (Agaricomycetes) in Mice with Nonalcoholic Steatohepatitis. Int J Med Mushrooms. 2020;22(7):683-692. [PMID: 32865925]
- Toshiaki Teratani, Kengo Tomita, Sachiko Toma-Fukai, Yutaro Nakamura, Toshimasa Itoh, Hikaru Shimizu, Yasunaga Shiraishi, Nao Sugihara, Masaaki Higashiyama, Takahiko Shimizu, Ikuo Inoue, Yasuhiro Takenaka, Ryota Hokari, Takeshi Adachi, Toshiyuki Shimizu, Soichiro Miura, Takanori Kanai. Redox-dependent PPAR γ /Tnpo1 complex formation enhances PPAR γ nuclear localization and signaling. Free Radic Biol Med. 2020 Aug 20;156:45-56. [PMID: 32553752]

Kahori Shimizu, Yuka Egusa, Syogo Nishimuta, Yuri Fukumura, Misato Yoshimura, Tomoya Inomoto, Tomoyuki Terada, Koji Tomita, Toru Nishinaka. Dietary calamondin supplementation slows the progression of non-alcoholic fatty liver disease in C57BL/6 mice fed a high-fat diet. *Int J Food Sci Nutr.* 2020 Aug 30;1-13. [PMID: 32862731]

Shinji Furuzono, Tetsuya Kubota, Junki Taura, Masahiro Konishi, Asuka Naito, Hiroshi Karasawa, Naoto Kubota, Takashi Kadowaki. A Novel Xanthene Derivative, DS20060511, Attenuates Glucose Intolerance by Inducing Skeletal Muscle-specific GLUT4 Translocation in Diabetic Mice. *Research Square (Preprint)* [PMID: none]

Yukiko Kinjo, Koji Wada, Moena Oe, De-Xing Hou, Makoto Takahashi. Effects of p-Hydroxybenzaldehyde and p-Hydroxyacetophenone from Non-centrifuged Cane Sugar, Kokuto, on Serum Corticosterone, and Liver Conditions in Chronically Stressed Mice Fed with a High-fat Diet. *Food Science and Technology Research.* Volume 26 (2020) Issue 4 [PMID: none]

Katsunori Nonogaki, Takao Kaji. Whey protein isolate inhibits hepatic FGF21 production, which precedes weight gain, hyperinsulinemia and hyperglycemia in mice fed a high-fat diet. *Sci Rep.* 2020 Sep 25;10(1):15784. [PMID: 32978487]

Kazuki Hattori, Hiroaki Wakatsuki, Chihiro Sakauchi, Shotaro Furutani, Sho Sugawara, Tomohisa Hatta, Tohru Natsume, Hidenori Ichijo. β -adrenergic receptor signaling evokes the PKA-ASK axis in mature brown adipocytes. *PLoS One.* 2020 Oct 27;15(10):e0232645. [PMID: 33108364]

Momoka Nishimura, Yuki Nomura, Moritoki Egi, Norihiko Obata, Makoto Tsunoda, Satoshi Mizobuchi. Suppression of behavioral activity and hippocampal noradrenaline caused by surgical stress in type 2 diabetes model mice. *BMC Neurosci.* 2020 Feb 17;21(1):8. [PMID: 32066381]

2021

Yukiomi Nakade, Rena Kitano, Kazumasa Sakamoto, Satoshi Kimoto, Taeko Yamauchi, Tadahisa Inoue, Yuji Kobayashi, Tomohiko Ohashi, Yoshio Sumida, Kiyoaki Ito, Masashi Yoneda. Characteristics of bile acid composition in high fat diet-induced nonalcoholic fatty liver disease in obese diabetic rats. *PLoS One.* 2021 Feb 24;16(2):e0247303. [PMID: 33626072]

Putra Santoso, Rita Maliza, Siti Jamalul Insani, Qonitah Fadhila, Resti Rahayu. Preventive effect of jicama (*Pachyrhizus erosus*) fiber against diabetes development in mice fed with high-fat diet. *Journal of Applied*

Pharmaceutical Science Vol. 11(01), pp 137-143, January, 2021. [PMID: none]

Gen Toshima, Takanobu Kiuchi, Yuka Iwama, Haruna Ohri, Fumiko Kimura, Shiho Nakagawa, Akira Sasaki, Kazuyuki Hiwatashi, Keishi Hata, Junichiro Takahashi. Anti-dyslipidemic effects of water extract from the leaves of *Petasites japonicus* subsp. *giganteus* (Akitabuki) *Journal of Biological Macromolecules* Volume 21 (2021) Issue 1 [PMID: none]

Koichi Fujisawa, Taro Takami, Haruko Shintani, Nanami Sasai, Toshihiko Matsumoto, Naoki Yamamoto, Isao Sakaida. Seasonal variations in photoperiod affect hepatic metabolism of medaka (*Oryzias latipes*) *FEBS Open Bio.* 2021 Jan 21. [PMID: 33475250]

(2021. 3)