

【糖尿病・肥満研究用高脂肪飼料Quick Fat参考文献リスト】

2005

遊佐寿恵, 片平清昭, 田口福志. GKラットにおける糖尿病発症促進飼料の給餌効果. 実験動物技術. 2005; 40(1): 7-14.

2006

Yamaguchi Y, Yamada K, Yoshikawa N, Nakamura K, Hagiwara J, Kunitomo M. Corosolic acid prevents oxidative stress, inflammation and hypertension in SHR/NDmc^r-cp rats, a model of metabolic syndrome. *Life Sci.* 2006; 79(26): 2474-9. [PMID: 16959274]

Matsumoto K, Watanabe Y, Ohya MA, Yokoyama S. Young persimmon fruits prevent the rise in plasma lipids in a diet-induced murine obesity model. *Biol Pharm Bull.* 2006; 29(12): 2532-5. [PMID: 17142997]

2007

Gao J, Katagiri H, Ishigaki Y, Yamada T, Ogiwara T, Imai J, Uno K, Hasegawa Y, Kanzaki M, Yamamoto TT, Ishibashi S, Oka Y. Involvement of apolipoprotein E in excess fat accumulation and insulin resistance. *Diabetes.* 2007; 56(1): 24-33. [PMID: 17192461]

Yoshizumi S, Suzuki S, Hirai M, Hinokio Y, Yamada T, Yamada T, Tsunoda U, Aburatani H, Yamaguchi K, Miyagi T, Oka Y. Increased hepatic expression of ganglioside-specific sialidase, NEU3, improves insulin sensitivity and glucose tolerance in mice. *Metabolism.* 2007; 56(3): 420-9. [PMID: 17292733]

Kanatani Y, Usui I, Ishizuka K, Bukhari A, Fujisaka S, Urakaze M, Haruta T, Kishimoto T, Naka T, Kobayashi M. Effects of pioglitazone on suppressor of cytokine signaling 3 expression: potential mechanisms for its effects on insulin sensitivity and adiponectin expression. *Diabetes.* 2007; 56(3): 795-803. [PMID: 17327450]

Chen KH, Mukaihoshi K, Sugihara H, Araki Y, Yamamoto G, Hattori T. High animal-fat intake changes the bile-acid composition of bile juice and enhances the development of Barrett's esophagus and esophageal adenocarcinoma in a rat duodenal-contents reflux model. *Cancer Sci.* 2007; 98(11): 1683-8. [PMID: 17868414]

Yanagiya T, Tanabe A, Iida A, Saito S, Sekine A, Takahashi A, Tsunoda T, Kamohara S, Nakata Y, Kotani K, Komatsu R, Itoh N, Mineo I, Wada J, Masuzaki H, Yoneda M, Nakajima A, Miyazaki S, Tokunaga K, Kawamoto M, Funahashi T, Hamaguchi K, Tanaka K, Yamada K,

Hanafusa T, Oikawa S, Yoshimatsu H, Nakao K, Sakata T, Matsuzawa Y, Kamatani N, Nakamura Y, Hotta K. Association of single-nucleotide polymorphisms in MTMR9 gene with obesity. *Hum Mol Genet.* 2007; 16(24): 3017-26. [PMID: 17855449]

2008

Tsuneki H, Murata S, Anzawa Y, Soeda Y, Tokai E, Wada T, Kimura I, Yanagisawa M, Sakurai T, Sasaoka T. Age-related insulin resistance in hypothalamus and peripheral tissues of orexin knockout mice. *Diabetologia.* 2008; 51(4): 657-67. [PMID: 18256806]

Ookawa K, Mochizuki K, Yokogoshi H. Effect of repeated stress in early childhood on the onset of diabetes mellitus in male Spontaneously Diabetic Torii rats. *J Vet Med Sci.* 2008; 70(2): 145-51. [PMID: 18319574]

Adachi T, Yanaka H, Kanai H, Nozaki M, Takahara Y, Tsuda M, Jonouchi T, Tsuda K, Hirasawa A, Tsujimoto G. Administration of perilla oil coated with Calshell increases glucagon-like peptide secretion. *Biol Pharm Bull.* 2008; 31(5): 1021-3. [PMID: 18451539]

Kunitomo M, Yamaguchi Y, Kagota S, Otsubo K. Beneficial effect of coenzyme Q10 on increased oxidative and nitritative stress and inflammation and individual metabolic components developing in a rat model of metabolic syndrome. *J Pharmacol Sci.* 2008; 107(2): 128-37. [PMID: 18544898]

Ijuin T, Yu YE, Mizutani K, Pao A, Tateya S, Tamori Y, Bradley A, Takenawa T. Increased insulin action in SKIP heterozygous knockout mice. *Mol Cell Biol.* 2008; 28(17): 5184-95. [PMID: 18573875]

Yamamoto Y, Ishino F, Kaneko-Ishino T, Shiura H, Uchio-Yamada K, Matsuda J, Suzuki O, Sato K. Type 2 diabetes mellitus in a non-obese mouse model induced by Meg1/Grb10 overexpression. *Exp Anim.* 2008; 57(4): 385-95. [PMID: 18633161]

Takayanagi Y, Matsumoto H, Nakata M, Mera T, Fukusumi S, Hinuma S, Ueta Y, Yada T, Leng G, Onaka T. Endogenous prolactin-releasing peptide regulates food intake in rodents. *J Clin Invest.* 2008; 118(12): 4014-24. [PMID: 19033670]

高橋 義宣, 河原崎 正貴, 星野 躍介, 本村 亜矢子, 江成 宏之. アンセリン含有サケエキスの高脂肪食飼育ラットに対する脂肪蓄積抑制効果. 日本水産学会誌, 2008; 74; 1075-81.

2009

Im R, Mano H, Matsuura T, Nakatani S, Shimizu J, Wada M. Mechanisms of blood glucose-lowering effect of aqueous extract from stems of Kothala himbutu (*Salacia reticulata*) in the mouse. *J Ethnopharmacol.* 2009; 121(2): 234-40. [PMID: 19028559]

Kunitomo M, Yamaguchi Y, Kagota S, Yoshikawa N, Nakamura K, Shinozuka K. Biochemical evidence of atherosclerosis progression mediated by increased oxidative stress in apolipoprotein E-deficient spontaneously hyperlipidemic mice exposed to chronic cigarette smoke. *J Pharmacol Sci.* 2009; 110(3): 354-61. [PMID: 19571461]

Beppu F, Niwano Y, Kyan R, Yasura K, Tamaki M, Nishino M, Midorikawa Y, Hamada H. Hypolipidemic Effects of Ethanol Extracts of *Citrus depressa* and *Annona atemoya*, Typical Plant Foodstuffs in Okinawa, Japan on KKAY Mice Fed with Moderately High Fat Diet. *Food Science and Technology Research* Vol. 15 (2009) No. 5 P 553-556 [No PMID]

2010

Mochiduki A, Takeda T, Kaga S, Inoue K. Stress response of prolactin-releasing peptide knockout mice as to glucocorticoid secretion. *J Neuroendocrinol.* 2010 Jun;22(6):576-84 [PMID: 20298457]

Takahashi M, Hori M, Kitahashi T, Imai T, Ishigamori R, Takasu S, Mutoh M, Sugimura T, and Wakabayashi K. Early and enhanced development of N-nitrosobis(2-oxopropyl)amine-induced hamster pancreatic adenocarcinomas by high-fat diet. *Cancer Res* April 15, 2010 70: 2478 [No PMID]

Watanabe S, Tsuneyama K. A triglyceride-lowering effect of cattle bile is associated with elevation of cholesterol levels and liver injury in mice. *Journal of Traditional Medicines* Vol. 27 (2010) No. 4 P 179-185 [No PMID]

Abe M, Toyohara T, Ishii A, Suzuki T, Noguchi N, Akiyama Y, Shiwaku HO, Nakagomi-Hagihara R, Zheng G, Shibata E, Souma T, Shindo T, Shima H, Takeuchi Y, Mishima E, Tanemoto M, Terasaki T, Onogawa T, Unno M, Ito S, Takasawa S, Abe T. The HMG-CoA reductase inhibitor pravastatin stimulates insulin secretion through organic anion transporter polypeptides. *Drug Metab Pharmacokinet.* 2010;25(3):274-82. [PMID: 20610886]

Mihara M, Aihara K, Ikeda Y, Yoshida S, Kinouchi M, Kurahashi K, Fujinaka Y, Akaike M, Matsumoto T. Inhibition of thrombin action ameliorates insulin resistance in type 2 diabetic db/db mice. *Endocrinology.* 2010

Feb;151(2):513-9 [PMID: 19966184]

2011

Doshi M, Takiue Y, Saito H, Hosoyamada M. The increased protein level of URAT1 was observed in obesity/metabolic syndrome model mice. *Nucleosides Nucleotides Nucleic Acids.* 2011 Dec;30(12):1290-4. [PMID: 22132989]

Fujisaka S, Usui I, Kanatani Y, Ikuuti M, Takasaki I, Tsuneyama K, Tabuchi Y, Bukhari A, Yamazaki Y, Suzuki H, Senda S, Aminuddin A, Nagai Y, Takatsu K, Kobayashi M, Tobe K. Telmisartan improves insulin resistance and modulates adipose tissue macrophage polarization in high-fat-fed mice. *Endocrinology.* 2011 May;152(5):1789-99. [PMID: 21427223]

Matsuda Y, Kobayashi M, Yamauchi R, Ojika M, Hiramitsu M, Inoue T, Katagiri T, Murai A, Horio F. Coffee and caffeine improve insulin sensitivity and glucose tolerance in C57BL/6J mice fed a high-fat diet. *Biosci Biotechnol Biochem.* 2011;75(12):2309-15. [PMID: 22146708]

Hashimoto M, Katakura M, Nabika T, Tanabe Y, Hossain S, Tsuchikura S, Shido O. Effects of hydrogen-rich water on abnormalities in a SHR.Cg-Leprcp/NDmcr rat - a metabolic syndrome rat model. *Med Gas Res.* 2011 Nov 3;1(1):26. [PMID: 22146083]

Hirata M, Suzuki M, Ishii R, Satow R, Uchida T, Kitazumi T, Sasaki T, Kitamura T, Yamaguchi H, Nakamura Y, Fukami K. Genetic defect in phospholipase C δ 1 protects mice from obesity by regulating thermogenesis and adipogenesis. *Diabetes.* 2011 Jul;60(7):1926-37. [PMID: 21617180]

2012

Takasaki M, Honma T, Yanaka M, Sato K, Shinohara N, Ito J, Tanaka Y, Tsuduki T, Ikeda I. Continuous intake of a high-fat diet beyond one generation promotes lipid accumulation in liver and white adipose tissue of female mice. *J Nutr Biochem.* 2012 Jun;23(6):640-5. [PMID: 21775120]

Kakino M, Sugiyama T, Kunieda H, Tazawa S, Maruyama H, Tsuruma K, Araki Y, Shimazawa M, Ichihara K, Mori H and Hara H. Agarwood (*Aquilaria Crassna*) Extracts Decrease High-protein Highfat Diet-induced Intestinal Putrefaction Toxins in Mice. *Pharm Anal Acta* 2012, 3:3 [No PMID]

Katakura M, Hashimoto M, Tanabe Y, Shido O. Hydrogen-rich water inhibits glucose and α,β -dicarbonyl compound-induced reactive oxygen species production in the SHR.Cg-Leprcp/NDmcr rat kidney. *Med Gas Res.*

- 2012 Jul 9;2(1):18. [PMID: 22776773]
- Hori M, Kitahashi T, Imai T, Ishigamori R, Takasu S, Mutoh M, Sugimura T, Wakabayashi K, Takahashi M. Enhancement of carcinogenesis and fatty infiltration in the pancreas in N-nitrosobis(2-oxopropyl)amine-treated hamsters by high-fat diet. *Pancreas*. 2011 Nov;40(8):1234-40. [PMID: 21989024]
- Honma T, Shinohara N, Ito J, Kijima R, Sugawara S, Arai T, Tsuduki T, Ikeda I. High-fat diet intake accelerates aging, increases expression of Hsd11b1, and promotes lipid accumulation in liver of SAMP10 mouse. *Biogerontology*. 2012 Apr;13(2):93-103. [PMID: 22033707]
- Takenaka T, Inoue T, Ohno Y, Miyazaki T, Nishiyama A, Ishii N, Suzuki H. Elucidating mechanisms underlying altered renal autoregulation in diabetes. *Am J Physiol Regul Integr Comp Physiol*. 2012 Sep 1;303(5):R495-504. [PMID: 22739351]
- Nagao M, Asai A, Kawahara M, Nakajima Y, Sato Y, Tanimura K, Okajima F, Takaya M, Sudo M, Takemitsu S, Harada T, Sugihara H, Oikawa S. Selective breeding of mice for different susceptibilities to high fat diet-induced glucose intolerance: Development of two novel mouse lines, Selectively bred Diet-induced Glucose intolerance-Prone and -Resistant. *J Diabetes Investig*. 2012 Jun 6;3(3):245-51. [PMID: 24843572]
- Mizukami H, Inaba W, Takahashi K, Inoue K, Sawanobori K, Yagihashi S. Augmented reduction of islet β -cell mass in Goto-Kakizaki rats fed high-fat diet and its suppression by pitavastatin treatment. *J Diabetes Investig*. 2012 Jun 6;3(3):235-44. [PMID: 24843571]
- Shanab AY, Nakazawa T, Ryu M, Tanaka Y, Himori N, Taguchi K, Yasuda M, Watanabe R, Takano J, Saido T, Minegishi N, Miyata T, Abe T, Yamamoto M. Metabolic stress response implicated in diabetic retinopathy: the role of calpain, and the therapeutic impact of calpain inhibitor. *Neurobiol Dis*. 2012 Dec;48(3):556-67. [PMID: 22967911]
- ## 2013
- Sugiishi A, Kimura M, Kamiya R, Ueki S, Yoneya M, Saito Y, Saito H. Derangement of ghrelin secretion after long-term high-fat diet feeding in rats. *Hepatol Res*. 2013 Oct;43(10):1105-14. [PMID: 23374505]
- Tsuduki T, Kitano Y, Honma T, Kijima R, Ikeda I. High dietary fat intake during lactation promotes development of diet-induced obesity in male offspring of mice. *J Nutr Sci Vitaminol (Tokyo)*. 2013;59(5):384-92. [PMID: 24418872]
- Nakaki J, Yamaguchi S, Torii Y, Inoue A, Minakami S, Kanno T, Murakami M, Tsuzuki M, Mochizuki H, Suyama K, Miyamoto M. Effect of fatty acids on the phosphate binding of TRK-390, a novel, highly selective phosphate-binding polymer. *Eur J Pharmacol*. 2013 Aug 15;714(1-3):312-7. [PMID: 23911881]
- Ishikawa M, Yoshida K, Okamura H, Ochiai K, Takamura H, Fujiwara N, Ozaki K. Oral *Porphyromonas gingivalis* translocates to the liver and regulates hepatic glycogen synthesis through the Akt/GSK-3 β signaling pathway. *Biochim Biophys Acta*. 2013 Dec;1832(12):2035-43. [PMID: 23899607]
- Ito R, Tsujihata Y, Matsuda-Nagasumi K, Mori I, Negoro N, Takeuchi K. TAK-875, a GPR40/FFAR1 agonist, in combination with metformin prevents progression of diabetes and β -cell dysfunction in Zucker diabetic fatty rats. *Br J Pharmacol*. 2013 Oct;170(3):568-80. [PMID: 23848179]
- ## 2014
- Hashiguchi C, Kawamoto S, Kasai T, Nishi Y, Nagaoka E. Influence of an antidiabetic drug on biomechanical and histological parameters around implants in type 2 diabetic rats. *Implant Dent*. 2014 Jun;23(3):264-9. [PMID: 24844386]
- Hattori T, Iizuka K, Horikawa Y, Takeda J. LRH-1 heterozygous knockout mice are prone to mild obesity. *Endocr J*. 2014;61(5):471-80. [PMID: 24531913]
- Koga S, Kojima A, Kuwabara S, Yoshiyama Y. Immunohistochemical analysis of tau phosphorylation and astroglial activation with enhanced leptin receptor expression in diet-induced obesity mouse hippocampus. *Neurosci Lett*. 2014 Jun 13;571:11-6. [PMID: 24785100]
- Nagao M, Asai A, Inaba W, Kawahara M, Shuto Y, Kobayashi S, Sanoyama D, Sugihara H, Yagihashi S, Oikawa S. Characterization of pancreatic islets in two selectively bred mouse lines with different susceptibilities to high-fat diet-induced glucose intolerance. *PLoS One*. 2014 Jan 13;9(1):e84725. [PMID: 24454742]
- Yonesaka T, Yoshida K, Iizuka S, Hagiwara H. Effects of fisetin on mouse lipid metabolism in vitro and in vivo. *Functional Foods in Health and Disease* 2014; 4(10):429-441 [No PMID]
- Hori H, Umezawa M, Uchiyama M, Niki R, Yanagita S, Takeda K. Effect of high-fat diet prior to pregnancy on hepatic gene expression and histology in mouse offspring. *J Perinat Med*. 2014 Jan;42(1):83-91 [PMID: 23974043]
- Imamura T, Ishizuka O, Ogawa T, Yamagishi T, Yokoyama H, Minagawa T, Nakazawa M, Gautam SS, Nishizawa O. Muscarinic receptors mediate cold stress-induced detrusor

overactivity in type 2 diabetes mellitus rats. *Int J Urol.* 2014 Oct;21(10):1051-8. [PMID: 24807830]

Koga S, Kojima A, Ishikawa C, Kuwabara S, Arai K, Yoshiyama Y. Effects of diet-induced obesity and voluntary exercise in a tauopathy mouse model: implications of persistent hyperleptinemia and enhanced astrocytic leptin receptor expression. *Neurobiol Dis.* 2014 Nov;71:180-92. [PMID: 25132556]

Suzuki S, Suzuki C, Hinokio Y, Ishigaki Y, Katagiri H, Kanzaki M, Azev VN, Chakraborty N, d'Alarcao M. Insulin-mimicking bioactivities of acylated inositol glycans in several mouse models of diabetes with or without obesity. *PLoS One.* 2014 Jun 27;9(6):e100466. [PMID: 24971987]

2015

Fushimi S, Miyazawa T, Nakagawa K, Burdeos GC, Miyazawa T. Young Persimmon Ingestion Suppresses Lipid Oxidation in Rats. *J Nutr Sci Vitaminol (Tokyo).* 2015;61(1):90-5. [PMID: 25994144]

Tsuduki T, Yamamoto K, E S1, Hatakeyama Y, Sakamoto Y. High Dietary Fat Intake during Lactation Promotes the Development of Social Stress-Induced Obesity in the Offspring of Mice. *Nutrients.* 2015 Jul 17;7(7):5916-32. [PMID: 26193313]

Imai N, Suzuki M, Hayashi K, Ishigami M, Hirooka Y, Abe T, Shioi G, Goto H, Fujimoto T. Hepatocyte-Specific Depletion of UBXD8 Induces Periportal Steatosis in Mice Fed a High-Fat Diet. *PLoS One.* 2015 May 13;10(5):e0127114. [PMID: 25970332]

Yamamoto K, E S1, Hatakeyama Y, Sakamoto Y, Tsuduki T. High-fat diet intake from senescence inhibits the attenuation of cell functions and the degeneration of villi with aging in the small intestine, and inhibits the attenuation of lipid absorption ability in SAMP8 mice. *J Clin Biochem Nutr.* 2015 Nov;57(3):204-11. [PMID: 26566305]

Shimizu N, Maruyama T, Yoshikawa N, Matsumiya R, Ma Y, Ito N, Tasaka Y, Kuribara-Souta A, Miyata K, Oike Y, Berger S, Schütz G, Takeda S, Tanaka H. A muscle-liver-fat signalling axis is essential for central control of adaptive adipose remodelling. *Nat Commun.* 2015 Apr 1;6:6693. [PMID: 25827749]

Nagao M, Asai A, Sugihara H, Oikawa S. Transgenerational changes of metabolic phenotypes in two selectively bred mouse colonies for different susceptibilities to diet-induced glucose intolerance. *Endocr J.* 2015;62(4):371-8. [PMID: 25736065]

Kishida T, Ejima A, Yamamoto K, Tanaka S, Yamamoto T3,

Mazda O. Reprogrammed Functional Brown Adipocytes Ameliorate Insulin Resistance and Dyslipidemia in Diet-Induced Obesity and Type 2 Diabetes. *Stem Cell Reports.* 2015 Oct 13;5(4):569-81. [PMID: 26365511]

Kato A, Naiki-Ito A, Nakazawa T, Hayashi K, Naitoh I, Miyabe K, Shimizu S, Kondo H, Nishi Y, Yoshida M, Umemura S, Hori Y, Mori T, Tsutsumi M, Kuno T, Suzuki S, Kato H, Ohara H, Joh T, Takahashi S. Chemopreventive effect of resveratrol and apocynin on pancreatic carcinogenesis via modulation of nuclear phosphorylated GSK3β and ERK1/2. *Oncotarget.* 2015 Dec 15;6(40):42963-75. [PMID: 26556864]

Kishida T, Ejima A, Yamamoto K, Tanaka S, Yamamoto T, Mazda O. Reprogrammed Functional Brown Adipocytes Ameliorate Insulin Resistance and Dyslipidemia in Diet-Induced Obesity and Type 2 Diabetes. *Stem Cell Reports.* 2015 Oct 13;5(4):569-81. [PMID: 26365511]

2016

Ohsaki Y, Kawai T, Yoshikawa Y, Cheng J, Jokitalo E, Fujimoto T. PML isoform II plays a critical role in nuclear lipid droplet formation. *J Cell Biol.* 2016 Jan 4;212(1):29-38. [PMID: 26728854]

Ito J, Nakagawa K, Kato S, Miyazawa T, Kimura F, Miyazawa T. The combination of maternal and offspring high-fat diets causes marked oxidative stress and development of metabolic syndrome in mouse offspring. *Life Sci.* 2016 Apr 15;151:70-5. [PMID: 26924496]

Li E, Nakata M, Shinozaki A, Yang Y, Zhang B, Yada T. Betatrophin expression is promoted in obese hyperinsulinemic type 2 but not type 1 diabetic mice. *Endocr J.* 2016 Apr 19. [Epub ahead of print] [PMID: 27097546]

Takemoto M, Takemoto H, Saijo R. Theaflavin Synthesized in a Selective, Domino-type, One-Pot Enzymatic Biotransformation Method with Camellia Sinensis Cell Culture Inhibits Weight Gain and Fat Accumulation to High-Fat Diet-Induced Obese mice. *Biol Pharm Bull.* 2016 May 25. [Epub ahead of print] [PMID: 27237789]

Kuda T, Masuko Y, Kawahara M, Kondo S, Nemoto M, Nakata T, Kataoka H, Takahashi H & Kimura B. Bile acid-lowering properties of *Lactobacillus plantarum* Sanriku-SU3 isolated from Japanese surfperch fish. *Food Bioscience.* 14, 41-46. [PMID: none]

Nakata M, Zhang B, Yang Y, Okada T, Shintani N, Hashimoto H, and Yada T. High-Fat Diet Augments VPAC1 Receptor-Mediated PACAP Action on the Liver, Inducing LAR Expression and Insulin Resistance. *Journal*

of Diabetes Research Volume 2016, Article ID 9321395, 10 pages. [PMID: 28044141]

Kitamura K, Takamura Y, Iwamoto T, Nomura M, Iwasaki H, Ohdera M, Murakoshi M, Sugiyama K, Matsuyama K, Manabe Y, Fujii NL, Fushiki T. Dammarane-type triterpene extracts of Panax notoginseng root ameliorates hyperglycemia and insulin sensitivity by enhancing glucose uptake in skeletal muscle. *Biosci Biotechnol Biochem.* 2016 Oct 24:1-8. [PMID: 27771987]

2017

Al Mamun A, Hashimoto M, Katakura M, Tanabe Y, Tsukichika S, Hossain S, Shido O. Effect of dietary n-3 fatty acids supplementation on fatty acid metabolism in atorvastatin-administered SHR.Cg-Leprcp/NDmc rats, a metabolic syndrome model. *Biomed Pharmacother.* 2017 Jan;85:372-379. [PMID: 27939244]

Nakata M, Yamamoto S, Okada T, Yada T. AAV-mediated IL-10 gene transfer counteracts inflammation in the hypothalamic arcuate nucleus and obesity induced by high-fat diet. *Neuropeptides.* 2017 Apr;62:87-92. [PMID: 27939689]

Kitamura K, Takamura Y, Iwamoto T, Nomura M, Iwasaki H, Ohdera M, Murakoshi M, Sugiyama K, Matsuyama K, Manabe Y, Fujii NL, Fushiki T. Dammarane-type triterpene extracts of Panax notoginseng root ameliorates hyperglycemia and insulin sensitivity by enhancing glucose uptake in skeletal muscle. *Biosci Biotechnol Biochem.* 2017 Feb;81(2):335-342. [PMID: 27771987]

Imai N, Suzuki M, Ishizu Y, Kuzuya T, Honda T, Hayashi K, Ishigami M, Hirooka Y, Ishikawa T, Goto H, Fujimoto T. Hepatocyte-specific depletion of ubiquitin regulatory X domain containing protein 8 accelerates fibrosis in a mouse non-alcoholic steatohepatitis model. *Histochem Cell Biol.* 2017 Sep;148(3):219-227. [PMID: 28421320]

2018

Minematsu A, Nishii Y, Sakata S. High-fat/high-sucrose diet results in higher bone mass in aged rats. *Bone Rep.* 2018 Jan 3;8:18-24. [PMID: 29322073]

Matsuzawa T, Yoshikawa T, Iida T, Kárpáti A, Kitano H, Harada R, Nakamura T, Sugawara A, Yamaguchi Y, Yanai K. Heparan sulfate in pancreatic β -cells contributes to normal glucose homeostasis by regulating insulin secretion. *Biochem Biophys Res Commun.* 2018 May 15;499(3):688-695. [PMID: 29605295]

Ohta T, Yamada T, Kamiya T, Gotoh T, Tsubaki, M, Shinohara M. Quick-fat diet inhibits the development of

diabetes in Spontaneously Diabetic Torii rats. *Thai Journal of Pharmaceutical Sciences* 2018, 42 (4): 183-187 [PMID: None]

山田久美子, 林櫻子, 飯沼光生 妊娠中のストレス下における咀嚼運動が出生後の仔の肥満に及ぼす影響について 岐阜歯科学会雑誌45巻1号、59-64、2018年7月 [PMID: None]

Ci Z, Jiang C, Feng S, Wu S, Cui Y, Sasaki Y and Kojima M. Anti-Obesity Effect of Proanthocyanidins from the Coat of Scarlet Runner Beans on High-Fat Diet-Fed Mice. *J of Food and Nutrition Research.* 2018, 6(2), 103-109 [PMID: None]

2019

Kanai S, Mukaisho KI, Yoshida S, Taniura N, Sugihara H. Host factors influence Barrett's carcinogenesis: findings from a mouse gastroduodenal reflux model. *Esophagus.* 2019 Jul;16(3):264-271. [PMID: 30790117]

2020

Akira Asai, Mototsugu Nagao, Koji Hayakawa, Teruo Miyazawa, Hitoshi Sugihara, Shinichi Oikawa. Leptin production capacity determines food intake and susceptibility to obesity-induced diabetes in Oikawa-Nagao Diabetes-Prone and Diabetes-Resistant mice. *Diabetologia.* 2020 Jun 19. [PMID: 32561946]

Mariko Seyama, Kaya Yoshida, Kayo Yoshida, Natsumi Fujiwara, Kisho Ono, Takanori Eguchi, Hotaka Kawai, Jiajie Guo, Yao Weng, Yuan Haoze, Kenta Uchibe, Mika Ikegame, Akira Sasaki, Hitoshi Nagatsuka, Kuniaki Okamoto, Hirohiko Okamura, Kazumi Ozaki. Outer membrane vesicles of *Porphyromonas gingivalis* attenuate insulin sensitivity by delivering gingipains to the liver. *Biochim Biophys Acta Mol Basis Dis.* 2020 Jun 1;1866(6):165731. [PMID: 32088316]

Dahlan Haslindah, Cangara Muhammad Husni, Masadah, Rina, Idris Irfan, Miskad Upik Anderiani, Zainuddin Andi Alfian, Mukani Miswani, Circulating and Renal Endothelin-1 Expression Lead to Glomerular Impairment in High Fat-Induced Diet Wistar Rats. *Indian Journal of Forensic Medicine & Toxicology*. Oct-Dec2020, Vol. 14 Issue 4, p8013-8018. [PMID: None]

Mariko Seyama, Kaya Yoshida, Kayo Yoshida, Natsumi Fujiwara, Kisho Ono, Takanori Eguchi, Hotaka Kawai, Jiajie Guo, Yao Weng, Yuan Haoze, Kenta Uchibe, Mika Ikegame, Akira Sasaki, Hitoshi Nagatsuka, Kuniaki Okamoto, Hirohiko Okamura, Kazumi Ozaki. Outer membrane vesicles of *Porphyromonas gingivalis* attenuate insulin sensitivity by delivering gingipains to the liver.

Biochim Biophys Acta Mol Basis Dis. 2020 Jun
1;1866(6):165731.[PMID: 32088316]

Yoshihiko Shibayama, Masanobu Nagano, Akira Fujii,
Kazunori Hashiguchi, Shunpei Morita, Yoshitada Kubo,
Tsutomu Nakagawa. Effect of concentrated Kurozu, a
traditional Japanese vinegar, on expression of hepatic
miR-34a, -149-3p, and -181a-5p in high-fat diet-fed mice.
Functional Foods in Health and Disease 2020; 10(1): 1-17.
[PMID: none]

2021

Hiroyuki Kato, Aya Naiki-Ito, Shugo Suzuki, Shingo
Inaguma, Masayuki Komura, Kenju Nakao, Taku Naiki,
Kenta Kachi, Akihisa Kato, Yoichi Matsuo, Satoru
Takahashi. DPYD, down-regulated by the potentially
chemopreventive agent luteolin, interacts with STAT3 in
pancreatic cancer. Carcinogenesis. 2021 Feb 26;bgab017.
[PMID: 33640964]

(2021.3)



日本クレア株式会社

<http://www.CLEA-Japan.com>

東京 A D 部 TEL. 03-5704-7050 FAX. 03-3792-2032
大阪 A D 部 TEL. 06-4861-7101 FAX. 06-4861-7108

仙台出張所 TEL. 022-352-4417 FAX. 022-352-4419
札幌出張所 TEL. 011-631-2725 FAX. 011-644-9209